

## Product datasheet for AM09005PU-N

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## C Reactive Protein (CRP) (19-224) Mouse Monoclonal Antibody [Clone ID: 5A9]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 5A9

**Applications:** ELISA, FC, IF, WB

Recommended Dilution: ELISA.

Western blot (1:1,000 - 1:2,000).

Immunocytochemistry/Immunoflourescence.

Flow cytometry.

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Recombinant human CRP (19-224 aa) purified from E. coli

**Specificity:** The antibody recognizes CRP at aa 19-224.

**Formulation:** PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

**Concentration:** lot specific

**Purification:** Protein-G affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** C-reactive protein, pentraxin-related

Database Link: Entrez Gene 1401 Human

P02741






Background:

C-reactive protein (CRP) is an acute-phase protein with a well-known association with infection and other inflammatory conditions. Mild elevations in C-reactive protein concentration predict myocardial infarction, stroke, and vascular death in a variety of clinical settings. Despite the lack of specific evidence that C-reactive protein levels are independently associated with cardiovascular risk in patients with hypertension, the prognostic value of C-reactive protein has proven to be complementary to that of blood pressure values.


Synonyms:

PTX1, C Reactive Protein, Pentraxin-related

## **Product images:**



Cell lysates of NIH-3T3, HepG2, Jurkat, HeLa and SH-SY5Y (each 40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CRP (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.



Lane1: NIH-3T3 Lane2: Hep G2
Lane3: Jurkat Lane4: HeLa

Lane5: SH-SY5Y

Western blot analysis: Cell lysates of NIH-3T3 (lane 1), HepG2 (lane 2), Jurkat (lane 3), HeLa (lane 4) and SH-SY5Y (lane 5) -each 40 ug-were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CRP (1:1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.