

Product datasheet for AM09055PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FLIP (CFLAR) (1-376) Mouse Monoclonal Antibody [Clone ID: AT8B12]

Product data:

Product Type: Primary Antibodies

Clone Name: AT8B12

Applications: ELISA, IF, WB

Recommended Dilution: ELISA.

Western Blot: 1/500-1/1000.

Immunofluorescence.

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant human FLIP (aa 1-376) purified from E. coli.

Specificity: The antibody recognizes FLIP.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CASP8 and FADD like apoptosis regulator

Database Link: Entrez Gene 12633 MouseEntrez Gene 117279 RatEntrez Gene 8837 Human

015519





Background: FLIP, also known as CASP8 and FADD-like apoptosis regulator (CFLAR), contains two death

effector domains (DEDs) and a caspase-like domain. FLIP may play a crucial role between cell survival and cell death pathway in mammalian cells and interacts with adapter protein FADD and Caspase-8 and -10, and potently inhibits apoptosis induced by all known death receptors DR3 (Death Receptor 3), TRAIL-R (TNF-related apoptosis-inducing ligand receptor) and TNFR1


(Tumor Necrosis Factor Receptor 1).

Synonyms: c-FLIP, CLARP, MRIT, CASH, FLAME-1, CASP8AP1, Usurpin

Protein Families: Druggable Genome, Protease

Protein Pathways: Apoptosis

Product images:



Western blot analysis: Cell lysates of mouse kidney, MCF7 and rat spleen (40 ug) were resolved by SDS-PAGE, trans-ferred to NC membrane and probed with FLIP antibody. (1/500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.