

Product datasheet for AM09140PU-L

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GAPDH Mouse Monoclonal Antibody [Clone ID: H8]

Product data:

Product Type: Primary Antibodies

Clone Name: H8

Recommended Dilution:

Applications: ELISA, WB

ELISA. Dot blot.

Western blott: 1/1,000-1/10,000 using ECL Substrate.

Reactivity: Bacteria, Human, Insect, Mouse, Rabbit, Rat, Yeast

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant GAPDH

Specificity: Recognizes native and recombinant GAPDH in Human, Mouse, Rat, Rabbit, sf9/nsect cells,

Yeast and BL-21 Bacteria.

Formulation: 0.01M PBS, pH 7.2

State: Purified

State: Lyophilized purified IgG fraction Preservative: 0.05% Sodium Azide

Reconstitution Method: Restore in double distilled or deioned H₂O to a concentration of 1.0 mg/ml.

Conjugation: Unconjugated

Storage: Store the antibody (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: One year from despatch.

Gene Name: glyceraldehyde-3-phosphate dehydrogenase

Database Link: Entrez Gene 14433 MouseEntrez Gene 24383 RatEntrez Gene 2597 Human

P04406





Background:

Glyceraldehyde 3 phosphate dehydrogenase (GAPDH) is well known as one of the key enzymes involved in glycolysis. As well as functioning as a glycolytic enzyme in cytoplasm, recent evidence suggests that mammalian GAPDH is also involved in a great number of intracellular proceses such as membrane fusion, microtubule bundling, phosphotransferase activity, nuclear RNA export, DNA replication, and DNA repair. During the last decade a lot of data appeared concerning the role of GAPDH in different pathologies including prostate cancer progression, programmed neuronal cell death, age related neuronal diseases, such as Alzheimer's and Huntington's disease. GAPDH is expressed in all cells. It is constitutively expressed in almost all tissues at high levels. There are however some physiological factors such as hypoxia and diabetes that increase GAPDH expression in certain cell types. GAPDH molecule is composed of four 36kDa subunits.

Synonyms: GAPD, CDABP0047

Protein Families: ES Cell Differentiation/IPS

Protein Pathways: Alzheimer's disease, Glycolysis / Gluconeogenesis, Metabolic pathways

Product images:

