

Product datasheet for AM09175PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL2 Mouse Monoclonal Antibody [Clone ID: 6B1]

Product data:

Product Type: Primary Antibodies

Clone Name: 6B1
Applications: ELISA

Recommended Dilution: ELISA: Clone 6B1 can match with clone 9F9 in Sandwich ELISA to detect the recombinant

human IL-2 antigen.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant Human Interleukin-2 (IL2)

Specificity: Recognizes Interleukin-2.

Formulation: 0.01M PBS, pH 7.2 without preservatives.

State: Aff - Purified

State: Lyophilized purified IgG fraction.

Reconstitution Method: Restore with double distillated water is recommended to adjust the final concentration to

1.00 mg/ml

Concentration: lot specific

Purification: Affinity Chromatography on Protein G.

Conjugation: Unconjugated

Storage: Store the antibody at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: interleukin 2

Database Link: Entrez Gene 3558 Human

P60568





IL2 Mouse Monoclonal Antibody [Clone ID: 6B1] - AM09175PU-N

Background: Interleukin 2 (IL2) is a secreted cytokine that is important for the proliferation of T and B

lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis like disease, which suggests an essential role of this gene in the immune response to antigenic stimuli. IL2 has been shown to have antitumor effects in

Synonyms: IL-2, TCGF

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Allograft rejection, Autoimmune thyroid disease, Cytokine-cytokine receptor interaction,

some studies. This is probably mediated by cytotoxic effector cells.

Graft-versus-host disease, Jak-STAT signaling pathway, T cell receptor signaling pathway, Type

I diabetes mellitus