

## **Product datasheet for AM09183HR-N**

## TNF alpha (TNF) Mouse Monoclonal Antibody [Clone ID: B1E4]

## **Product data:**

**Product Type:** Primary Antibodies

Clone Name: B1E4
Applications: ELISA
Recommended Dilution: ELISA.
Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Purified recombinant human TNF-α

**Specificity:** This monoclonal antibody recognizes human TNF-α.

**Formulation:** 50% 0.01M pH7.2 PBS 50% glycerol

Label: HRP

State: Liquid purified Ig

Label: Horseradish Peroxidase

**Purification:** Affinity chromatography on Protein G

Conjugation: HRP

Storage: Store the antibody at -20°C.


Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** tumor necrosis factor

Database Link: Entrez Gene 7124 Human

P01375



9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Background:

Tumor Necrosis Factor Alpha (TNF alpha) is a protein secreted by lipopolysaccharide stimulated macrophages, and causes tumor necrosis when injected into tumour bearing mice. TNF alpha is believed to mediate pathogenic shock and tissue injury associated with endotoxemia. TNF alpha exists as a multimer of two, three, or five noncovalently linked units, but shows a single 17 kDa band following SDS PAGE under non reducing conditions. TNF alpha is closely related to the 25 kDa protein Tumour Necrosis Factor beta (lymphotoxin), sharing the same receptors and cellular actions. TNF alpha causes cytolysis or cytostasis of certain transformed cells, being synergistic with interferon gamma in its cytotoxicity. Although it has little effect on many cultured normal human cells, TNF alpha appears to be directly toxic to vascular endothelial cells. Other actions of TNF alpha include stimulating growth of human fibroblasts and other cell lines, activating polymorphonuclear neutrophils and osteoclasts, and induction of interleukin 1, prostaglandin E2 and collagenase production. TNF alpha is currently being evaluated in treatment of certain cancers and AIDS Related Complex.

**Synonyms:** TNF, TNF-a, TNFA, TNFSF2, Cachectin

**Protein Families:** Druggable Genome, Secreted Protein, Transcription Factors, Transmembrane

**Protein Pathways:** Adipocytokine signaling pathway, Allograft rejection, Alzheimer's disease, Amyotrophic lateral

sclerosis (ALS), Apoptosis, Asthma, Cytokine-cytokine receptor interaction, Dilated cardiomyopathy, Fc epsilon RI signaling pathway, Graft-versus-host disease, Hematopoietic cell lineage, Hypertrophic cardiomyopathy (HCM), MAPK signaling pathway, Natural killer cell mediated cytotoxicity, NOD-like receptor signaling pathway, RIG-I-like receptor signaling pathway, Systemic lupus erythematosus, T cell receptor signaling pathway, TGF-beta signaling

pathway, Toll-like receptor signaling pathway, Type I diabetes mellitus, Type II diabetes

mellitus