

## Product datasheet for AM09224PU-N

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Helicobacter pylori (CagA Protein) Mouse Monoclonal Antibody [Clone ID: 3C10]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 3C10
Applications: ELISA

**Recommended Dilution: ELISA:** Helicobacter pylori antibody (Clone 3C10) can be used as Detection antibody.

In Sandwich ELISA, anti-HP CagA antibody clone 10E9 (Cat.-No AM09221PU-N) and anti-HP CagA antibody clone 5H10 (Cat.-No AM09222PU-N) coated wells match with HRP conjugated

clone 3C10 to detect the recombinant antigen.

HRP conjugated 3C10 also selectively detected cell lysate of a CagA containing HP strain when

clone 10E9 was used as coating antibody.

**Reactivity:** Helicobacter pylori

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: A highly immunogenic 32Kd CagA fragment, recombinant.

**Specificity:** This antibody is raised against the *H. pylori* cytotoxin associated gene A protein (CagA).

Reacts with the 32Kd recombinant Helicobacter pylori CagA. Cross reaction with other proteins

has not been identified.

**Formulation:** 0.01M PBS, pH 7.0 without preservatives

State: Aff - Purified

State: Lyophilized purified IgG fraction

**Reconstitution Method:** Restore with Double distillated water to adjust the final concentration to 1.0 mg/ml.

**Purification:** Affinity Chromatography on Protein G

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.





## Helicobacter pylori (CagA Protein) Mouse Monoclonal Antibody [Clone ID: 3C10] - AM09224PU-N

Background: Helicobacter pylori (H. pylori), a spiral rod shaped gram-negative bacterium, is frequently found

in the stomach. In infected populations, 10-20% may develop gastritis and gastric ulcer and 1-2% may develop cancer. The genome of *H. pylori* isolates from carriers with symptoms contains a 40kb pathogenicity island encoding *H. pylori cytotoxin*, cytotoxin associated gene A

protein (CagA) and other virulence associated factors. CagA is used as a biomarker for

virulent *H. pylori* strains.

Synonyms: H. pylori