

Product datasheet for AM09281PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD41-CD61 Complex Mouse Monoclonal Antibody [Clone ID: 8]

Product data:

Product Type: Primary Antibodies

Clone Name: 8

Applications: ELISA, WB

Recommended Dilution: This GPIIb/IIIa antibody is Suitable for use in ELISA and Western Blot.

Use of this Monoclonal antibody at a concentration of 10-20 µg/ml (culture supernatants) will

allow visualization of NP 40 solubilized human normal platelets.

The titer of purified ascites was determined by ELISA coated with 100 µl of 1x10e8 washed

platelets.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: 10e9 Human platelets suspension.

The supernatant fluid from positive clones was screened by ELISA coated with washed

platelets.

Specificity: Human platelet membrane glycoproteins IIb/IIIa (GP IIb/IIIa).

Under reducing conditions, this antibody will bind against the Human platelet membrane glycoproteins IIb/IIIa complex, which have a molecular weights of ~130,000 (IIb) and 115,000

(IIIa).

Formulation: 0.01M PBS, pH 7.2 without preservatives.

State: Aff - Purified

State: Lyophilized purified IgG fraction.

Reconstitution Method: Restore with Double distillated water to adjust the final concentration to 1.0 mg/ml.

Purification: Affinity Chromatography on Protein G.

Conjugation: Unconjugated

Storage: Store the antibody at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.





Synonyms:

CD41-CD61 Complex Mouse Monoclonal Antibody [Clone ID: 8] - AM09281PU-N

Background: The integrins are a family of more than 23 heterodimeric transmembrane proteins that

mediate cell-cell adhesions as well as cell-substratum adhesions and signal transduction processes. Integrins are heterodimers consisting of noncovalently associated alpha and beta subunits. CD61 non-covalently associates with CD41 (alpha Ilb integrin) and is expressed by megakaryocytes and platelets. The CD61/CD41 complex acts as a receptor for such adhesive ligands as fibronectin, fibrinogen and von Willebrand factor during platelet stimulation.

GPIIb-IIIa, ITGA2B, GP2B, ITGAB, Platelet membrane glycoprotein IIb, GPalpha IIb, GPIIb,

ligarius as fibroffectiff, fibriffogeri and vort Willebrand factor during plateiet stiffulation.

GP3A, CD41, CD61, Integrin alpha-IIb, Integrin beta-3, ITGB3