

## **Product datasheet for AM09316PU-N**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ELK1 pSer383 Mouse Monoclonal Antibody [Clone ID: D7H2]**

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: D7H2
Applications: ELISA

**Recommended Dilution:** Indirect ELISA: the antibody is reactive to the un-conjugated Serine383 phosphorylated

human ELK-1 peptide, and reactive to the serine383 phosphorylated human ELK-1 peptide

conjugated with KLH or the peptide-BSA conjugate.

The suggested antigen Coating quantity is 1 µg/ml of peptide/or peptide conjugating in pH

9.5 carbonate buffer, 100 µl/well.

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Peptide of a phosphorylated fragment of ELK1 (pSer383) conjugated with KLH.

Specificity: Reactive to the un-conjugated serine 383 phosphorylated Human ELK-1 peptide fragment, and

reactive to the Serine383 phosphorylated Human ELK-1 peptide fragment conjugated to KLH

or the peptide- BSA conjugate. Not reactive to KLH or BSA.

**Formulation:** 0.01M PBS, pH 7.0 without preservatives

State: Aff - Purified

State: Lyophilized purified IgG fraction

**Reconstitution Method:** Restore with Double distillated water to adjust the final concentration to 1.0 mg/ml.

**Purification:** Affinity Chromatography on Protein G

Conjugation: Unconjugated

Storage: Upon receipt, store (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** ELK1, ETS transcription factor

Database Link: Entrez Gene 2002 Human

P19419





## ELK1 pSer383 Mouse Monoclonal Antibody [Clone ID: D7H2] - AM09316PU-N

**Background:** ELK1 belongs to the EST transcription factor family. Proteins in the family have a unique DNA

binding domain that binds to conserved DNA sequences. Phosphorylation of ELK1 at serine 383 and 389 by mitogen-activated protein kinases (MAPKs) activates ELK1. The binding of ELK1 to target sequences is also regulated by its co-factor, serum response factor (SRF), by alternative splicing, and by SUMOylation. ELK1 regulates the transcription of c-fos and genes encoding cytoskeleton proteins, and was found to play a role in longterm memory formation,

drug addiction and other neurological disorders.

Synonyms: Elk-1

**Protein Families:** Druggable Genome, Transcription Factors

**Protein Pathways:** Endometrial cancer, ErbB signaling pathway, Focal adhesion, GnRH signaling pathway, Insulin

signaling pathway, MAPK signaling pathway, Prion diseases