

Product datasheet for AM09325PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

STAT5B pTyr694 Mouse Monoclonal Antibody [Clone ID: A2H6]

Product data:

Product Type: Primary Antibodies

Clone Name: A2H6
Applications: ELISA

Recommended Dilution: Indirect ELISA: the antibody is reactive to Tyrosine699 phosphorlylated Human STAT5b

peptide fragment conjugated with KLH or the peptide-BSA conjugate.

The suggested antigen coating quantity is 1 ug/ml of peptide/or peptide conjugating in pH 9.5

carbonate buffer, 100 ul/well.

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: A Tyrosine 699 phosphorylated Human STAT5b peptide fragment conjugated with KLH.

Specificity: Reactive to Tyrosine 699 phosphorlylated human STAT5b peptide fragment conjugated with

KLH or the peptide-BSA conjugate, but not reactive to unconjugated KLH or BSA.

Formulation: 0.01M PBS, pH 7.2 without preservatives

State: Aff - Purified

State: Lyophilized purified IgG fraction

Reconstitution Method: Restore with Double distillated water to adjust the final concentration to 1.0 mg/ml.

Purification: Affinity chromatography on Protein G

Conjugation: Unconjugated

Storage: Upon receipt, store (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: signal transducer and activator of transcription 5B

Database Link: Entrez Gene 6777 Human

P51692





STAT5B pTyr694 Mouse Monoclonal Antibody [Clone ID: A2H6] - AM09325PU-N

Background: Stat5b is a transcription factor that is activated in response to extra-cellular stimuli including

IL-2, IL-4, MCSF and a variety of growth factors. In the process, receptor associated kinases phosphorylate Stat5b. Stat5b, then, forms dimers and translocates to nucleus. Inside nucleus, Stat5b binds to the promoter region on various genes to regulate biological processes such as apoptosis, T cell receptor signalling, and adult mammary gland development. Fusion of Stat5b to retinoic acid receptor-alpha (RARα) gene was identified in a small subgroup of acute

promyelocytic leukemia patients.

Synonyms: STAT-5B

Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Stem cell relevant signaling - JAK/STAT

signaling pathway, Transcription Factors

Protein Pathways: Acute myeloid leukemia, Chemokine signaling pathway, Chronic myeloid leukemia, ErbB

signaling pathway, Jak-STAT signaling pathway, Pathways in cancer