

Product datasheet for AM09361PU-S

CLEC4E Mouse Monoclonal Antibody [Clone ID: AT16E3]

Product data:

Product Type:	Primary Antibodies
Clone Name:	AT16E3
Applications:	ELISA, IF, IHC, WB
Recommended Dilution:	ELISA. Western blot: 1/250-1/1000. Immunofluorescence: 1/250-1/500. Immunohistochemistry on Paraffin Embedded tissues: 1/50.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Recombinant Human MINCLE (41-219aa) purified from E. coli
Specificity:	The antibody recognizes Human CLEC4E (MINCLE). Other species not tested.
Formulation:	PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol State: Purified State: Liquid purified Ig fraction
Concentration:	lot specific
Purification:	Protein-G Affinity Chromatography
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	C-type lectin domain family 4 member E
Database Link:	<u>Entrez Gene 26253 Human</u> <u>Q9ULY5</u>



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CLEC4E Mouse Monoclonal Antibody [Clone ID: AT16E3] – AM09361PU-S

Background: MINCLE (macrophage inducible C-type lectin, also called CLEC4E or CLECSF9), which is a diverse family of protein which was originally defined by their ability to recognize a wide range of ligand of carbohydrate structure. MINCLE expressed in macrophages subjected to several types of stress. It plays an essential role in response to trehalose-6,6'-dimycolate (TDM) and activated by a synthetic analogue, trehalose dibehenate (TDB). Recently it was reported that MINCLE is associated with an immunoreceptor tyrosine-based activation motif-containing Fc receptor γ chain (FcRγ) and functions as an activating receptor for damaged self- and non-self-pathogenic fungi.

Protein Families:

Druggable Genome, Transmembrane

Product images:



Cell lysates of HeLa (35ug) were resolved by SDS-PAGE, transferred to NC membrane and probed with anti-human MINCLE (1/1000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.



Paraffin embedded sections of colorectal cancer tissue were incubated with anti-Human MINCLE antibody (1/50) for 2 hours at room temperature. Antigen retrieval was performed in 0.1M sodium citrate buffer and detected using Diaminobenzidine (DAB).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US





Immunofluorescence of human HeLa cells stained with monoclonal anti-human MINCLE antibody (1/500) with Texas Red (Red). Nucleus was stained by Hoechst 33342 (Blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US