

Product datasheet for AM09387PU-N

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

CTBP1 Mouse Monoclonal Antibody [Clone ID: AT4D6]

Product data:

Product Type: Primary Antibodies

Clone Name: AT4D6

Applications: ELISA, FC, IF, WB

Recommended Dilution: ELISA.

Western blot (1:500). Immunofluorescence. Flow Cytometry.

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Recombinant human CTBP1 (1-440aa) purified from E. coli

Specificity: The antibody recognizes human CTBP1.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-G affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: C-terminal binding protein 1

Database Link: Entrez Gene 1487 Human

Q13363





Background: The C-terminal-binding protein (CtBP) was originally identified as a cellular protein that

> interacts with a C-terminal motif of adenovirus E1A oncoproteins. CtBP is highly conserved in higher eukaryotes. Vertebrates contain two highly homologous CtBP proteins (CtBP1 and CtBP2) while invertebrates contain a single CtBP. CtBP family proteins share significant sequence homology with D-2-hydroxy acid dehydrogenases. Also, CtBP proteins associate


with various proteins and functions as a co-repressor of transcription.

C-terminal-binding protein 1, CTBP Synonyms:


Protein Pathways: Chronic myeloid leukemia, Notch signaling pathway, Pathways in cancer, Wnt signaling

pathway

Product images:



The cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CTBP1 (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.



Western blot: Cell lysates (40ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human CTBP1 (1:500). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.