

Product datasheet for **AM09391PU-S**

NRAS (1-186) Mouse Monoclonal Antibody [Clone ID: AT2G9]

Product data:

Product Type:	Primary Antibodies
Clone Name:	AT2G9
Applications:	ELISA, FC, IF, WB
Recommended Dilution:	ELISA. Western blot (1:500 - 1:5000; recommended starting dilution is 1:1000). Immunocytochemistry / Immunofluorescence. Flow cytometry.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Recombinant human NRAS (1-186aa) purified from <i>E. coli</i>
Specificity:	The antibody recognizes human RAS at aa 1-186. Other species not tested.
Formulation:	PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol State: Purified State: Liquid purified Ig fraction
Concentration:	lot specific
Purification:	Protein-G affinity chromatography
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	neuroblastoma RAS viral oncogene homolog
Database Link:	Entrez Gene 4893 Human P01111



[View online »](#)



Background: The Ras protein is a subfamily of small GTPases that are involved in signal transduction. They are activated by ligand-binding receptor tyrosine kinases such as EGFR, PDGFR, CSF and FGF. These kinases transiently convert RAS-GDP (inactive form) to RAS-GTP, the active form of RAS. The main genes are HRAS, NRAS and KRAS.

Synonyms: NRAS, HRAS1, KRAS, KRAS2, RASK2, HRAS, HRAS1



Protein Families: Druggable Genome

Protein Pathways: Acute myeloid leukemia, Axon guidance, B cell receptor signaling pathway, Bladder cancer, Chemokine signaling pathway, Chronic myeloid leukemia, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Gap junction, Glioma, GnRH signaling pathway, Insulin signaling pathway, Long-term depression, Long-term potentiation, MAPK signaling pathway, Melanogenesis, Melanoma, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pathways in cancer, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, T cell receptor signaling pathway, Thyroid cancer, Tight junction, VEGF signaling pathway

Product images:



The extracts of 293T (35ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human RAS (1:500~1:5000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. The cross-reacting of anti-human RAS (AT2G9) was analyzed by western blot with recombinant protein (200ng) of NRAS, KRAS and HRAS.



Western blot analysis: The extracts of 293T (35ug) were resolved by SDS-PAGE, transferred to PVDF membrane and probed with anti-human RAS (1:500-1:5000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system. The cross-reacting of anti-human RAS (AT2G9) was analyzed by Western blot with recombinant protein (200ng) of NRAS, KRAS and HRAS.