

Product datasheet for AM10019PU-S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SERPINA12 Mouse Monoclonal Antibody [Clone ID: 1C4]

Product data:

Product Type: Primary Antibodies

Clone Name: 1C4

Applications: ELISA, WB **Recommended Dilution:** ELISA.

Western blot (1:1,000 - 1:3,000), recommended starting dilution is 1:2000.

Reactivity: Human, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant human Vaspin (21-414 aa) purified from E. coli

Specificity: The antibody recognizes human and rat Vaspin.

Other species not tested.

Formulation: PBS, pH 7.4 containing 0.02% Sodium Azide and 10% Glycerol

State: Purified

State: Liquid purified Ig fraction

Concentration: lot specific

Purification: Protein-G affinity chromatography

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C for up to two weeks or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: serpin family A member 12

Database Link: Entrez Gene 145264 Human

Q8IW75



SERPINA12 Mouse Monoclonal Antibody [Clone ID: 1C4] - AM10019PU-S


Background:

Vaspin (visceral adipose-specific SERPIN), a newly identified adipokine, which is a member of serine protease inhibitor family. Vaspin is also a unique insulin sensitizing adipocytokine in obesity. A recent publication indicates that induction of human vaspin mRNA expression in adipose tissue is regulated in a fat depot-specific manner and could be associated with parameters of obesity, insulin resistance, and glucose metabolism.


Synonyms:

Serpin A12, Vaspin, OL-64

Product images:



Rat serum (15ug) was resolved by SDS-PAGE, transferred to NC membrane and probed with anti-human Vaspin (1:2000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.



Western blot analysis: Rat serum (15 ug) was resolved by SDS-PAGE, transferred to NC membrane and probed with anti-human Vaspin (1:2000). Proteins were visualized using a goat anti-mouse secondary antibody conjugated to HRP and an ECL detection system.