

## Product datasheet for **AM10036FC-N**

### LOC104968698 Mouse Monoclonal Antibody [Clone ID: S100217]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	S100217
Applications:	FC, IF, IHC
Recommended Dilution:	<b>Immunohistochemistry:</b> 1/50-1/100, use streptavidin~biotin system or polymer system, incubate 30 minutes at room temperature. FFPE tissue section requires antigen retriever (boiling tissue in 10 mM citrate, pH 6.0 for 20 mins, followed by cooling at RT for 20 mins). <b>Immunocytochemistry.</b> <b>Immunofluorescence:</b> 10-20 µg/ml (1/10-1/20), incubate for 2 hours in the dark at RT or it can also be incubated overnight at 4°C. <b>Flow Cytometry:</b> 0.2-1.0 µg/0.1 ml (1/200-1/1,000). <i>Recommended Positive Control:</i> Melanoma or schwannoma.
Reactivity:	Bovine, Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Purified bovine S100 protein
Specificity:	This antibody stains schwannoma, ependymomas, astroglomas, almost all benign and malignant melanoma and their metasis. Cellular Localization: Cytoplasmic.
Formulation:	PBS, pH 7.4 Label: FITC State: Liquid purified Ig fraction. Stabilizer: 1% BSA Preservative: 0.05% Sodium Azide
Concentration:	lot specific
Purification:	Protein A Chromatography.
Conjugation:	FITC



[View online »](#)

<b>Storage:</b>	Store the antibody undiluted at 2-8°C. This product is photosensitive and should be protected from light.
<b>Stability:</b>	Shelf life: One year from despatch.
<b>Database Link:</b>	<a href="#">Entrez Gene 528735 Bovine P02639</a>
<b>Background:</b>	<p>S100 protein is a type of low molecular weight protein found in vertebrates characterized by two calcium binding sites of the helix-loop-helix ("EF-hand type") conformation. There are at least 21 different types of S100 proteins. The name is derived from the fact that the protein is 100% Soluble in ammonium sulfate at neutral pH. Most S100 proteins are homodimeric, consisting of two identical polypeptides held together by non-covalent bonds. S100A is composed of an alpha and beta chain whereas S100B is composed of two beta chains. Although S100 proteins are structurally similar to calmodulin, they differ in that they are cell-specific, expressed in particular cells at different levels depending on environmental factors. To contrast, calmodulin is a ubiquitous and universal intracellular Ca<sup>++</sup> receptor widely expressed in many cells. S-100 is normally present in cells derived from the neural crest (Schwann cells, melanocytes, and glia cells), chondrocyte, adipocytes, myoepithelial cells, macrophages, Langerhans cells, dendritic cells, and keratinocytes. It may be present in some breast epithelial cells.</p>
<b>Synonyms:</b>	S100 Calcium binding protein, S-100 protein, S100A, S100B, S-100 protein alpha chain, S-100 protein beta chain, Astrocyte Marker