

Product datasheet for **AM10093SU-N**

Hepatocyte Specific Antigen (HSA) Mouse Monoclonal Antibody [Clone ID: OCH1E5]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OCH1E5
Applications:	IHC
Recommended Dilution:	Immunohistochemistry on Formalin-Fixed, Paraffin-Embedded Sections: 1/100. Pretreatment of deparaffinized tissue with heat-induced epitope retrieval is recommended. Use Polymer anti Mouse/Rabbit IgG as a detection system. Positive Control: Hepatocellular carcinoma.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Formalin fixed failed Human allograft liver that was mechanically disrupted actual price paid for the product.
Specificity:	Clone OCH1E5 is useful in studying hepatocellular tumors. It may be useful in differentiating clear cell hepatocellular carcinomas from other clear cell malignancies. It has been shown in the literature to be useful in differentiating hepatoblastoma of embryonal type from small round cell tumors of childhood. OCH1E5 labels an antigen in the mitochondrial fraction of the liver homogenates. Cellular Localization: Cytoplasmic.
Formulation:	State: Supernatant State: Tissue Culture Supernatant with 0.2% BSA and 15mM Sodium Azide
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Background:	Hepatoblastoma is the most common primary tumor of the liver in children. The use of specific hepatocyte markers and also of alpha Fetoprotein or carcinoembryonic antigen are useful for the identification of normal and malignant fetal hepatocytes.
Synonyms:	hepatocellular carcinoma marker, hepatocyte marker, HepPar-1



[View online »](#)

Product images:



Formalin-Fixed, Paraffin-Embedded Human liver stained with HepPar-1 antibody Cat.-No. AM10093SU-N using peroxidase conjugate and DAB chromogen. Note the cytoplasmic staining of hepatocytes.