

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM11001PU-N

MET pTyr1234/1235 Mouse Monoclonal Antibody [Clone ID: 6AT1877]

Product data:

Product Type:	Primary Antibodies
Clone Name:	6AT1877
Applications:	WB
Recommended Dilution:	ELISA: 1/1000. Western Blot.
Reactivity:	Human, Mouse
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	KLH conjugated synthetic peptide corresponding to sequences in Tyr1234/1235 region of human Met and residues both Tyr1234 and Tyr1235 are phosphorylated.
Specificity:	This antibody detects MET/HGFR pTyr1234/1235.
Formulation:	PBS with 0.09% (W/V) Sodium Azide as preservative State: Purified State: Liquid purified lg fraction. Preservative: 0.09% (W/V) Sodium Azide
Concentration:	lot specific
Purification:	Protein G Chromatography eluted with high and low pH buffers and neutralized immediately, followed by dialysis against PBS.
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	MET proto-oncogene, receptor tyrosine kinase
Database Link:	<u>Entrez Gene 4233 Human</u> <u>P08581</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

CRIGENE MET pTyr1234/1235 Mouse Monoclonal Antibody [Clone ID: 6AT1877] – AM11001PU-N

Background:MET/HGFR is a receptor for hepatocyte growth factor, with tyrosine-protein kinase activity.
The MET/HGFR Type I membrane protein contains 1 Sema domain and consists of a
heterodimer formed of an alpha chain (50 kDa) and a beta chain (145 kDa) which are disulfide
linked. In the fully processed c-Met product, the alpha subunit is extracellular, and the beta
subunit has extracellular, transmembrane, and tyrosine kinase domains as well as sites of
tyrosine phosphorylation. Two isoforms for the protein have been described. Activation of
MET after rearrangement with the TPR gene produces an oncogenic protein. MET is
overexpressed in a significant percentage of human cancers and is amplified during the
transition between primary tumors and metastasis. Defects in MET are a cause of hereditary
papillary renal carcinoma (HPRC), also known as papillary renal cell carcinoma 2 (RCCP2).
HPRC is a form of inherited kidney cancer characterized by a predisposition to develop
multiple, bilateral papillary renal tumors. The pattern of inheritance is consistent with
autosomal dominant transmission with reduced penetrance.

Synonyms: Hepatocyte growth factor receptor, MET, Scatter factor receptor, HGF/SF receptor, c-Met

Product images:



Figure 1. Detection of endogenous Met in HepG2 cell line. 10 ug/lane of HepG2 cell lysate was used to examine the expression of human Met. Lanes 1-5 represent different anti-Met monoclonal antibodies that are [AM11002PU-N], [AM11003PU-N], [AM11004PU-N], [AM11005PU-N] and [AM11006PU-N]. Lane 6 represents autophosohorylated-Met in HepG2 cell line detected by anti-phospho-Met Mab ().

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US