

Product datasheet for AM11036PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SUV39H1 (N-term) Mouse Monoclonal Antibody [Clone ID: 42AT239.96.72]

Product data:

Product Type: Primary Antibodies

Clone Name: 42AT239.96.72

Applications: WB

Recommended Dilution: ELISA: 1/1,000.

Western blotting: 1/100-1/500.

Reactivity: Human, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: This antibody was raised using purified recombinant GST fusion protein encoding N-terminal

of human SUV39H1.

Specificity: This antibody is specific to SUV39H1.

Formulation: PBS containing 0.09% (W/V) Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein G Chromatography eluted with high and low pH buffers and neutralized immediately,

followed by dialysis against PBS.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: suppressor of variegation 3-9 homolog 1

Database Link: Entrez Gene 20937 MouseEntrez Gene 6839 Human

<u>043463</u>





Background:

Similar to acetylation and phosphorylation, histone methylation at the N-terminal tail has emerged as an important role in regulating chromatin dynamics and gene activity. Histone methylation occurs on arginine and lysine residues and is catalyzed by two families of proteins, the protein arginine methyltransferase family and the SET-domain-containing methyltransferase family. Five members have been identified in the arginine methyltransferase family. About 27 are grouped into the SET-domain family, and another 17 make up the PR domain family that is related to the SET domain family. The retinoblastoma protein-interacting zinc finger geneRIZ1 is a tumor suppressor gene and a FOUNDING member of the PR domain family. RIZ1 inactivation is commonly found in many types of human cancers and occurs through loss of mRNA expression, frame shift mutation, chromosomal deletion, and missense mutation. RIZ1 is also a tumor susceptibility gene in mice. The loss of RIZ1 mRNA in human cancers was shown to associate with DNA methylation of its promoter CpG island. Methylation of the RIZ1 promoter strongly correlated with lost or decreased RIZ1 mRNA expression in breast, liver, colon, and lung cancer cell lines as well as in liver cancer tissues.

Synonyms: SUV39H, Lysine N-methyltransferase 1A

Protein Families: Druggable Genome
Protein Pathways: Lysine degradation

Product images:




Figure 1. Western analysis of extracts from 293 cells using SUV39H1 antibody.