

Product datasheet for AM12004PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cd247 Hamster Monoclonal Antibody [Clone ID: H146-968]

Product data:

Product Type: Primary Antibodies

Clone Name: H146-968

Applications: ELISA, FC, IF, WB

Recommended Dilution: This clone (H146-968) has been reported to work in Flow Cytometry (in permeabilized cells),

Western Blotting and ELISA.

Reactivity: Human, Mouse

Host: Hamster

Isotype: IgG

Clonality: Monoclonal

Immunogen: Synthetic peptide KLH conjugated corresponding to amino acids 151-164 of Mouse CD3 zeta

Specificity: This antibody detects the zeta subunit of the T cell receptor (TCR).

Formulation: PBS containing 0.02% Sodium Azide as preservative.

State: Purified

State: Liquid purified Ig fraction.

Concentration: lot specific

Purification: Protein G Affinity Chromatography.

Conjugation: Unconjugated

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Gene Name: CD247 antigen

Database Link: Entrez Gene 919 HumanEntrez Gene 12503 Mouse

P24161



Cd247 Hamster Monoclonal Antibody [Clone ID: H146-968] - AM12004PU-N

Background:

The TCR complex consists of a ligand specific alpha/beta heterodimer non-covalently associated with five invariant chains including the CD3 gamma, delta, eta and zeta subunits. This complex regulates assembly and expression of the receptor and is thought to be responsible for transmembrane transduction of signals after binding to TCR $\alpha\beta$. Studies have suggested that the CD3z subunit (also know as CD247) plays an important role in two distinct cellular compartments. In the cytoplasm, it may function to regulate Ag receptor expression and in the plasma membrane it may be required for optimal signaling by the physiologic ligand Ag/MHC. The CD3z subunit is also associated with the CD16 IgG Fc receptor on NK cells.

Synonyms: T3Z, TCRZ, CD3 zeta chain

Product images:



Western Blot Analysis of CD3z antibody: Lysate from Mouse Spleen and Thymus (Thymus shown) cells probed with at 1.0 ug/ml. AP conjugated Goat Anti-Hamster IgG used as a secondary antibody.