

Product datasheet for AM12006PU-N

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Rockville, MD 20850, US

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

CD28 Mouse Monoclonal Antibody [Clone ID: CD28.2]

Product data:

Product Type: Primary Antibodies

Clone Name: CD28.2

Applications: FC, FN, IF, IHC, IP, WB

Recommended Dilution: Flow cytometry.

Immunoprecipitation.

Western blot.

Immunohistochemistry (frozen sections).

Immunocytochemistry.

Functional application: T cell costimulation.

Reactivity: Human, Primate

lgG1

Host: Mouse

Isotype:

Monoclonal Clonality:

DC28.1.3.3 murine T cell hybridoma transfected with human CD28 cDNA Immunogen:

This antibody reacts with CD28, a disulfide-linked homodimeric type I glycoprotein (monomer Specificity:

of Mw 44 kDa) which is a critical costimulatory receptor of T cells.

Formulation: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4

State: Purified

State: Liquid Ig fraction

Concentration: lot specific

Purification: Protein A affinity chromatography; purity > 95% (by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

DO NOT FREEZE!

Stability: Shelf life: one year from despatch.

Gene Name: CD28 molecule

Entrez Gene 940 Human Database Link:

P10747





Background: CD28 is the critical T cell costimulatory receptor which provides to the cell the important

second activation signal by binding CD80 and CD86 that are expressed by antigen presenting cells. Besides its costimulation role CD28 functions in preventing T cells from anergic hyporesponsive state or from undergoing premature apoptotic cell death. CD28 is also

expressed on human fetal NK cells and some NK cell lines, whereas on murine NK cells the

CD28 expression is much broader.

Synonyms: TP44


Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Allograft rejection, Autoimmune thyroid disease, Cell adhesion molecules (CAMs), Graft-

versus-host disease, Systemic lupus erythematosus, T cell receptor signaling pathway, Type I

diabetes mellitus, Viral myocarditis

Product images:



Surface staining of human peripheral blood leukocytes with anti-human CD28 (CD28.2) purified.