

## Product datasheet for **AM20602PU-N**

### CDC6 Mouse Monoclonal Antibody [Clone ID: IMD-6]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	IMD-6
Applications:	WB
Recommended Dilution:	Western Blot: 1 - 2 µg/ml.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Recombinant human Cdc6
Specificity:	This antibody reacts to CDC6.
Formulation:	1.2 % sodium acetate, with 2 mg BSA and 0.01 mg sodium azide as preservative. State: Aff - Purified State: Lyophilized purified Ig fraction
Reconstitution Method:	Restore with 1.2% sodium acetate or neutral PBS
Concentration:	0,1 mg/ml (after reconstitution with PBS)
Purification:	Affinity chromatography
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at -20°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	cell division cycle 6
Database Link:	<a href="#">Entrez Gene 990 Human Q99741</a>



[View online »](#)

<b>Background:</b>	Cell division cycle(CDC25) gene product is a protein-tyrosine phosphatase, activates a partially purified p34(cdc2)/cyclin B complex. The cdc25 protein also shares homology with a protein phosphatase with activity against both tyrosine and serine (and thus probably threonine) phosphate residues. CDC25C is expressed predominantly in G2 phase in healer cells. The human gene encodes a protein with a predicted molecular mass of 53,000 daltons whose C-terminal domain shares about 37% sequence identity with the fission yeast cdc25 mitotic inducer. CDC25C gene is mapped to 5q31.
<b>Synonyms:</b>	CDC18L, CDC6-related protein, p62(cdc6), HsCDC6, Cdc18-related protein, HsCdc18
<b>Protein Pathways:</b>	Cell cycle