

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM26006AC-N

CD19 Mouse Monoclonal Antibody [Clone ID: 4G7]

Product data:


Product Type:	Primary Antibodies
Clone Name:	4G7
Applications:	FC
Recommended Dilution:	Flow Cytometry analysis of human blood cells using 10 μ l reagent / 100 μ l of whole blood or 10 ⁶ cells in a suspension.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Human CCL (chronic lymphocytic leukemia) cells
Specificity:	This antibody recognizes CD19 (B4), a 95 kDa type I transmembrane glycoprotein of immunoglobulin superfamily, expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.
Formulation:	Phosphate buffered saline (PBS) containing 15 mM sodium azide Label: APC State: Liquid purified lg fraction Label: Conjugated with cross-linked Allophycocyanin under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use.
Conjugation:	APC
Storage:	Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be protected from light.
Stability:	Shelf life: one year from despatch.
Gene Name:	CD19 molecule
Database Link:	<u>Entrez Gene 930 Human</u> <u>P15391</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CD19 Mouse Monoclonal Antibody [Clone ID: 4G7] – AM26006AC-N
Background:	CD19 is a transmembrane glycoprotein of Ig superfamily expressed by B cells from the time of heavy chain rearrangement until plasma cell differentiation. It forms a tetrameric complex with CD21 (complement receptor type 2), CD81 (TAPA-1) and Leu13. Together with BCR (B cell antigen receptor), this complex signals to decrease B cell treshold for activation by the antigen. Besides being signal-amplifying coreceptor for BCR, CD19 can also signal independently of BCR coligation and it turns out to be a central regulatory component upon which multiple signaling pathways converge. Mutation of the CD19 gene results in hypogammaglobulinemia, whereas CD19 overexpression causes B cell hyperactivity.
Synonyms:	Leu-12, B-cell marker
Protein Families	Druggable Genome, Transmembrane
Protein Pathway	s: B cell receptor signaling pathway, Hematopoietic cell lineage, Primary immunodeficiency

Product images:



Surface staining of human peripheral blood leukocytes with anti-human CD19 (4G7) APC.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US