

Product datasheet for **AM26006BT-N**

CD19 Mouse Monoclonal Antibody [Clone ID: 4G7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	4G7
Applications:	FC
Recommended Dilution:	Indirect immunofluorescence analysis by Flow Cytometry .
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human CCL (chronic lymphocytic leukemia) cells
Specificity:	The mouse monoclonal antibody 4G7 recognizes CD19 (B4), a 95 kDa type I transmembrane glycoprotein of immunoglobulin superfamily, expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.
Formulation:	PBS Label: Biotin State: Liquid purified Ig fraction Preservative: 15 mM sodium azide Label: Conjugated with Biotin-LC-NHS under optimum conditions. The reagent is free of unconjugated biotin.
Concentration:	lot specific
Conjugation:	Biotin
Storage:	Store at 2-8°C. Do not freeze.
Stability:	Shelf life: One year from despatch.
Gene Name:	CD19 molecule
Database Link:	Entrez Gene 930 Human P15391




[View online »](#)

Background:

CD19 is a transmembrane glycoprotein of Ig superfamily expressed by B cells from the time of heavy chain rearrangement until plasma cell differentiation. It forms a tetrameric complex with CD21 (complement receptor type 2), CD81 (TAPA-1) and Leu13. Together with BCR (B cell antigen receptor), this complex signals to decrease B cell threshold for activation by the antigen. Besides being signal-amplifying coreceptor for BCR, CD19 can also signal independently of BCR coligation and it turns out to be a central regulatory component upon which multiple signaling pathways converge. Mutation of the CD19 gene results in hypogammaglobulinemia, whereas CD19 overexpression causes B cell hyperactivity.

Synonyms:

Leu-12, B-cell marker

Product images:

Surface staining of human peripheral blood leukocytes with anti-human CD19 (4G7) purified.