

Product datasheet for **AM26006FC-N**

CD19 Mouse Monoclonal Antibody [Clone ID: 4G7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	4G7
Applications:	FC
Recommended Dilution:	Flow Cytometry analysis of human blood cells using 20 µl reagent / 100 µl of whole blood or 10 ⁶ cells in a suspension.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human CCL (chronic lymphocytic leukemia) cells
Specificity:	This antibody recognizes CD19 (B4), a 95 kDa type I transmembrane glycoprotein of immunoglobulin superfamily, expressed on B lymphocytes and follicular dendritic cells; it is lost on plasma cells.
Formulation:	Phosphate buffered saline (PBS) containing 15 mM sodium azide Label: FITC State: Liquid purified Ig fraction Label: Conjugated with Fluorescein isothiocyanate under optimum conditions. The reagent is free of unconjugated and adjusted for direct use.
Conjugation:	FITC
Storage:	Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be protected from light.
Stability:	Shelf life: one year from despatch.
Gene Name:	CD19 molecule
Database Link:	Entrez Gene 930 Human P15391



[View online »](#)

Background:

CD19 is a transmembrane glycoprotein of Ig superfamily expressed by B cells from the time of heavy chain rearrangement until plasma cell differentiation. It forms a tetrameric complex with CD21 (complement receptor type 2), CD81 (TAPA-1) and Leu13. Together with BCR (B cell antigen receptor), this complex signals to decrease B cell threshold for activation by the antigen. Besides being signal-amplifying coreceptor for BCR, CD19 can also signal independently of BCR coligation and it turns out to be a central regulatory component upon which multiple signaling pathways converge. Mutation of the CD19 gene results in hypogammaglobulinemia, whereas CD19 overexpression causes B cell hyperactivity.

Synonyms:


Leu-12, B-cell marker

Protein Families:

Druggable Genome, Transmembrane

Protein Pathways:

B cell receptor signaling pathway, Hematopoietic cell lineage, Primary immunodeficiency

Product images:

Surface staining of human peripheral blood leukocytes with anti-human CD19 (4G7) purified.