

## Product datasheet for AM26006RD5-N

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD19 Mouse Monoclonal Antibody [Clone ID: 4G7]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: 4G7
Applications: FC

Recommended Dilution: Flow Cytometry analysis of human blood cells using 4 µl reagent / 100 µl of whole blood or

106 cells in a suspension.

The content of a vial (0.4 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human CCL (chronic lymphocytic leukemia) cells

Specificity: This antibody recognizes CD19 (B4), a 95 kDa type I transmembrane glycoprotein of

immunoglobulin superfamily, expressed on B lymphocytes and follicular dendritic cells; it is

lost on plasma cells.

**Formulation:** Phosphate buffered saline (PBS)

Label: PE-DL594

State: Liquid purified lg fraction Preservative: 15 mM sodium azide

Label: Conjugated with tandem dye PE-DyLight™ 594 (PE-DL594) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use.

Conjugation: PE-DL594

Storage: Store undiluted at 2-8°C. DO NOT FREEZE! This products is photosensitive and should be

protected from light.

**Stability:** Shelf life: one year from despatch.

Gene Name: CD19 molecule

Database Link: Entrez Gene 930 Human

P15391





## CD19 Mouse Monoclonal Antibody [Clone ID: 4G7] - AM26006RD5-N

**Background:** CD19 is a transmembrane glycoprotein of Ig superfamily expressed by B cells from the time

of heavy chain rearrangement until plasma cell differentiation. It forms a tetrameric complex with CD21 (complement receptor type 2), CD81 (TAPA-1) and Leu13. Together with BCR (B cell

antigen receptor), this complex signals to decrease B cell treshold for activation by the antigen. Besides being signal-amplifying coreceptor for BCR, CD19 can also signal

independently of BCR coligation and it turns out to be a central regulatory component upon

which multiple signaling pathways converge. Mutation of the CD19 gene results in hypogammaglobulinemia, whereas CD19 overexpression causes B cell hyperactivity.

Synonyms: Leu-12, B-cell marker

**Note:** DyLight<sup>™</sup> is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.