

Product datasheet for AM26020AC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD62P (SELP) Mouse Monoclonal Antibody [Clone ID: AK4]

Product data:

Product Type: Primary Antibodies

Clone Name: AK4
Applications: FC

Recommended Dilution: Flow Cytometry analysis of human blood cells using 10 μl reagent / 100 μl of whole blood or

106 cells in a suspension.

Reactivity: Human, Primate

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human platelets

Specificity: The antibody recognizes CD62P (P-selectin), a 140 kD single chain type I transmembrane

glycoprotein present in secretory alpha-granules in platelets, in Weibel-Palade bodies in endothelial cells and in megakaryocytes; it is relocated to the plasma membrane upon

activation.

Formulation: Phosphate buffered saline (PBS)

Label: APC

State: Liquid Ig fraction

Stabilizer: 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA)

Preservative: 15 mM sodium azide

Label: Conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No

reconstitution is necessary.

Conjugation: APC

Storage: Store undiluted at 2-8°C. DO NOT FREEZE! This products is photosensitive and should be

protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: selectin P

Database Link: Entrez Gene 6403 Human

P16109






Background:

CD62P (P-selectin) is an adhesion glycoprotein that is expressed on platelets and endothelial cells upon their activation. Interaction between CD62P and its mucin-like ligand PSGL-1 (P-selectin glycoprotein ligand-1) expressed on the microvilli of most leukocytes supports leukocyte rolling along postkapillary venules at the earliest time of inflammation. Both CD62P and PSGL-1 are extended glycoproteins that form homodimers. CD62P dimerization is probably mediated through interactions of the transmembrane domains and stabilizes leukocyte tethering and rolling, probably by increasing rebinding within a bond cluster.

Synonyms:

SELP, GMRP, GRMP, PADGEM, GMP-140, LECAM3

Product images:



Surface staining of human peripheral blood with anti-CD62P (AK4) APC.