

Product datasheet for AM26029FC-N

Cd4 Rat Monoclonal Antibody [Clone ID: GK1.5]

Product data:

Product Type: Primary Antibodies

Clone Name: GK1.5

Applications: FC

Recommended Dilution: Flow cytometry: $3 \mu g/ml$.

Reactivity: Mouse **Host:** Rat

Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Mouse CTL clone V4 cells

Specificity: This antibody reacts with an extracellular epitope of mouse CD4 transmembrane

glycoprotein (55 kDa).

Formulation: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.4

Label: FITC

State: Liquid purified Ig fraction

Label: Conjugated with Fluorescein isothiocyanate (FITC) under optimum conditions. The

reagent is free of unconjugated FITC.

Concentration: lot specific

Conjugation: FITC

Storage: Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be

protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: CD4 antigen

Database Link: Entrez Gene 12504 Mouse

P06332



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Background:

CD4 is a single chain transmembrane glycoprotein of immunoglobulin supergene family. In its extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like C2-type). The intracellular region of CD4 associates with p56Lck, a Src-like protein tyrosine kinase. It was described that CD4 segregates into specific detergent-resistant T-cell membrane microdomains. CD4 binds to MHC class II molecules (by CDR2-like region in CD4 domain 1), HIV envelope protein gp120 (by CDR2-like region in CD4 domain 1) and other ligands, such as IL-16 (by to CD4 domain 3) or L-selectin. CD4 is a co-receptor involved in immune response (co-receptor activity in binding to MHC class II molecules) and HIV infection. CD4 regulates T-cell activation, T/B-cell adhesion, T-cell diferentiation, T-cell selection and signal transduction. Defects in antigen presentation (MHC class II) cause dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and organs (SCID immunodeficiency).


Synonyms: T-cell surface antigen T4/Leu-3

Product images:




Surface staining of CD4 in murine splenocytes with anti-CD4 (GK1.5) APC.





Surface staining of CD4 in murine splenocytes with anti-CD4 (GK1.5) azide free, DAR/APC.



Surface staining of CD4 in murine splenocytes with anti-CD4 (GK1.5) PE.