

9620 Medical Pochvillo, MD

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

Product datasheet for AM26029RP-N

Cd4 Rat Monoclonal Antibody [Clone ID: GK1.5]

Product data:

Product Type:	Primary Antibodies
Clone Name:	GK1.5
Applications:	FC
Recommended Dilution:	Flow cytometry: 3 µg/ml.
Reactivity:	Mouse
Host:	Rat
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Mouse CTL clone V4 cells
Specificity:	This antibody reacts with an extracellular epitope of mouse CD4 transmembrane glycoprotein (55 kDa).
Formulation:	PBS, approx. pH 7.4 Label: PE State: Liquid purified Ig fraction Preservative: 15 mM sodium azide Label: Conjugated with R-Phycoerythrin under optimum conditions. The conjugate is purified by size-exclusion chromatography.
Concentration:	lot specific
Conjugation:	PE
Storage:	Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be protected from light.
Stability:	Shelf life: one year from despatch.
Gene Name:	CD4 antigen
Database Link:	<u>Entrez Gene 12504 Mouse</u> <u>P06332</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Cd4 Rat Monoclonal Antibody [Clone ID: GK1.5] – AM26029RP-N

Background:CD4 is a single chain transmembrane glycoprotein of immunoglobulin supergene family. In
its extracellular region there are 4 immunoglobulin-like domains (1 Ig-like V-type and 3 Ig-like
C2-type). The intracellular region of CD4 associates with p56Lck, a Src-like protein tyrosine
kinase. It was described that CD4 segregates into specific detergent-resistant T-cell
membrane microdomains. CD4 binds to MHC class II molecules (by CDR2-like region in CD4
domain 1), HIV envelope protein gp120 (by CDR2-like region in CD4 domain 1) and other
ligands, such as IL-16 (by to CD4 domain 3) or L-selectin. CD4 is a co-receptor involved in
immune response (co-receptor activity in binding to MHC class II molecules) and HIV
infection. CD4 regulates T-cell activation, T/B-cell adhesion, T-cell diferentiation, T-cell
selection and signal transduction. Defects in antigen presentation (MHC class II) cause
dysfunction of CD4+ T-cells and their almost complete absence in patients blood, tissue and
organs (SCID immunodeficiency).

Synonyms:

T-cell surface antigen T4/Leu-3

Product images:



Surface staining of CD4 in murine splenocytes with anti-CD4 (GK1.5) PE.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US