

## Product datasheet for AM26053PU-N

# **OriGene Technologies, Inc.**9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## CD4 Mouse Monoclonal Antibody [Clone ID: EDU-2]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: EDU-2
Applications: FC, IHC

**Recommended Dilution:** Immunohistochemistry on frozen sections: 0.2 μg/ml (1:1000).

Has been described to work in FACS.

Suggested positive control: Human tonsil.

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Stimulated human leukocytes
Specificity: This antibody detects CD4.

Formulation: Stock solution contains 0.2 mg/ml lgG, phosphate buffered saline pH 7.2 (PBS), 5 mg/ml

bovine serum albumin (BSA), and 0.09% sodium azide as a preservative.

State: Aff - Purified

State: Lyophilized Ig fraction

Label: Reconstitute by adding 0.5ml distilled water.

**Purification:** Affinity chromatography

Conjugation: Unconjugated

**Storage:** Prior to reconstitution store at 2-8°C. Following reconstitution store (in aliquots) at -20°C.

Avoid repeated freezing and thawing.

**Stability:** Shelf life: one year from despatch.

Gene Name: Homo sapiens CD4 molecule (CD4), transcript variant 1

Database Link: Entrez Gene 920 Human

P01730



### CD4 Mouse Monoclonal Antibody [Clone ID: EDU-2] - AM26053PU-N

**Background:** CD4 is a glycoprotein expressed on the surface of T helper cells, regulatory T cells,

monocytes, macrophages, and dendritic cells. It is mainly expressed by the T lymphocyte subset that recognizes antigens associated with self MHC class II molecules. CD4 is the primary receptor for HIV retroviruses. Like many cell surface markers, it is a member of the

immunoglobulin superfamily.

**Synonyms:** T-cell surface antigen T4/Leu-3

**Note:** The antigen is CD4, a 55kD glycoprotein (reduced and nonreduced). The epitope has not

been further characterized.

Antigen distribution:

Isolated cells: the antibody stains approximately 20-60% of human peripheral blood mononuclear cells in flow cytometry.


Protein Families: Adult stem cells, Druggable Genome, ES Cell Differentiation/IPS, Induced pluripotent stem

cells, Transmembrane

**Protein Pathways:** Antigen processing and presentation, Cell adhesion molecules (CAMs), Hematopoietic cell

lineage, Primary immunodeficiency, T cell receptor signaling pathway

## **Product images:**



Immunohistochemistry on frozen sections: Staining with AM26053PU-N on human tonsil