

Product datasheet for AM26067RP-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

STRO-1 Mouse Monoclonal Antibody [Clone ID: STRO-1]

Product data:

Product Type: Primary Antibodies

Clone Name: STRO-1

Applications: FC

Recommended Dilution: Flow cytometry.

Reactivity: Human
Host: Mouse
Isotype: IgM

Clonality: Monoclonal

Immunogen: Human CD34 positive bone marrow cells

Specificity: This antibody recognizes the cell surface antigen STRO-1 expressed by bone marrow

mesenchymal stromal cells and nucleated erythroid precursors, but not by committed

hematopoietic progenitors.

Formulation: Tris buffered saline (TBS)

Label: PE

State: Liquid purified Ig fraction

Stabilizer: 0.2% (w/v) high-grade protease free Bovine Serum Albumin (BSA)

Preservative: 15 mM sodium azide

Label: Conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is

purified by size-exclusion chromatography.

Concentration: lot specific

Conjugation: PE

Storage: Store the antibody at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Stability: Shelf life: one year from despatch.






Background:

STRO-1 is a cell surface antigen expressed by stromal elements in human bone marrow, identified by monoclonal antibody STRO-1. Approximately 10% of mononuclear cells, greater than 95% of which are nucleated erythroid precursors, are STRO-1 positive, whereas the CFU-GM (colony-forming unit granulocyte-macrophage), BFU-E (erythroid burst) and CFU-Mix (mixed colonies) committed progenitor cells are negative. CFU-F (fibroblast colony-forming cells) are present exclusively in the STRO-1 positive population. When plated under long-term bone marrow culture conditions, STRO-1 positive cells generate adherent cell layers containing multiple stromal cell types, including adipocytes, smooth muscle cells, osteoblasts, chondrocytes, and fibroblastic elements. In combination with glycophorin A, STRO-1 is a useful marker for identification of mesenchymal stem cells. STRO-1 and CD117 are markers for osteosarcoma cells.

Product images:



Surface staining of human peripheral blood with anti-STRO-1 (STRO-1) PE.