

Product datasheet for AM26138RP-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD163 Mouse Monoclonal Antibody [Clone ID: GHI/61]

Product data:

Product Type: Primary Antibodies

Clone Name: GHI/61

Applications: FC

Recommended Dilution: Flow cytometry analysis of human blood cells using 10 μl reagent / 100 μl of whole blood or

10e6 cells in a suspension.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Hairy cell leukemia cells

Specificity: This antibody recognizes CD163, an approximately 130 kDa high affinity scavenger receptor

expressed mainly on monocytes and macrophages, which binds hemoglobin-haptoglobin

complex.

Formulation: Phosphate buffered saline (PBS) containing 15 mM sodium azide and 0.2% (w/v) high-grade

protease free Bovine Serum Albumin (BSA) as a stabilizing agent

Label: PE

State: Liquid purified Ig fraction

Purification: Conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is purified by

size-exclusion chromatography and adjusted for direct use.

Conjugation: PE

Storage: Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be

protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: CD163 molecule

Database Link: Entrez Gene 9332 Human

O86VB7






Background:

CD163, also known as M130, is a member of the scavenger receptor family, accounting for the clearance of hemoglobin-haptoglobin complexes during limited hemolysis, which protects the body, in particular the kidneys, against heme-mediated oxidative damages. It does not have measurable affinity for noncomplexed hemoglobin or haptoglobin. Immunomodulatory role of CD163 has been postulated. CD163 is expressed by cells of the monocyte-macrophage lineage and its extracellular part also circulates in plasma as a soluble protein, especially during sepsis and other conditions affecting macrophage activity, when its level may raise manyfold.

Synonyms: M130, Hemoglobin scavenger receptor, Macrophage marker

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Product images:



Surface staining of human peripheral blood using anti-human CD163 (clone GHI/61) PE.