

## Product datasheet for AM26142RP-N

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Integrin alpha E (ITGAE) Mouse Monoclonal Antibody [Clone ID: Ber-ACT8]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: Ber-ACT8

Applications: FC

**Recommended Dilution:** Flow Cytometry analysis of human blood cells using 10 μl reagent / 100 μl of whole blood or

10<sup>6</sup> cells in a suspension.

The content of a vial (1 ml) is sufficient for 100 tests.

Reactivity: Human, Primate

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** HTLV-1 induced human T cell line MAPS16

Specificity: The mouse monoclonal antibody Ber-ACT8 recognizes CD103 (alpha E integrin), a type I

transmembrane glycoprotein primarily found on intestinal intraepithelial lymphocytes.

Formulation: PBS

Label: PE

State: Liquid purified Ig fraction

Stabilizer: 0.2% (w/v) high-grade protease free Bovine Serum Albumin

Preservative: 15 mM sodium azide

Label: Conjugated with R-Phycoerythrin (PE) under optimum conditions. The conjugate is

purified by size-exclusion chromatography and adjusted for direct use.

Conjugation: PE

Storage: Store the antibody at 2 - 8 °C. DO NOT FREEZE! This product is photosensitive and should be

protected from light.

**Stability:** Shelf life: one year from despatch.

Gene Name: integrin subunit alpha E

Database Link: Entrez Gene 3682 Human

P38570





## Integrin alpha E (ITGAE) Mouse Monoclonal Antibody [Clone ID: Ber-ACT8] - AM26142RP-N

Background: CD103 / Integrin alphaE is an integrin subunit that is expressed on intraepithelial

lymphocytes, epithelial dendritic cells, lamina propria-derived dendritic cells, a subpopulation of lamina propria T cells, a small subset of peripheral lymphocytes, namely T reg cells, and on activated and TGF-beta stimulated lymphocytes. CD103 is in mature form cleaved into two disulfide-linked chains (C-terminal 150 kDa chain and N-terminal 25 kDa chain). It heterodimerizes with integrin beta7 subunit to form alphaE/beta7 integrin (mucosal

lymphocyte 1 antigen), which through binding E-cadherin mediates homing of lymphocytes to the intestinal epithelium, and, in addition to the role in adhesion, may serve as an accessory

molecule for intraepithelial lymphocyte activation.

Synonyms: Integrin alpha-E, Integrin alpha-IEL, HML-1 antigen, ITGAE