

## Product datasheet for **AM26182PU-N**

### **TREM1 Mouse Monoclonal Antibody [Clone ID: 6B1]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	6B1
Applications:	ELISA, FN, WB
Recommended Dilution:	Flow cytometry (Typical starting working dilution is 1:50. Functional assays. Immunoassays. Western blot (Typical starting working dilution is 1:50).
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Recombinant TREM-1 - IgG1 fusion protein
Specificity:	This antibody recognizes human triggering receptor expressed on myeloid cells 1 (TREM-1). It recognizes both membrane bound and soluble TREM-1. Also, the antibody neutralizes the interaction between TREM-1 and its ligand.
Formulation:	PBS State: Purified State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 0.1% bovine serum albumin
Concentration:	lot specific
Purification:	Protein G
Conjugation:	Unconjugated
Storage:	Store at 2 - 8 °C.
Stability:	Shelf life: one year from despatch.
Gene Name:	triggering receptor expressed on myeloid cells 1
Database Link:	<a href="#">Entrez Gene 54210 Human Q9NP99</a>



[View online »](#)

**Background:**

TREM-1 is a transmembrane glycoprotein belonging to a family related to the natural killer cell receptors. TREM-1 is an important activating receptor, of 26 kDa, involved in the innate inflammatory response and in sepsis. TREM-1 is expressed at low levels in the early development of the hematopoietic system and in the promonocytic stage and at high levels on neutrophil granulocytes (PMN), monocytes, and macrophage subsets. The expression of TREM-1 is upregulated by microbial products, that is, by toll-like receptor ligands such as lipoteichoic acid (LTA) of Gram-positive or lipopolysaccharide (LPS) of Gram-negative bacteria. Ligation of TREM-1 is synergistic with TLR agonists on the activation of receptor bearing cells. Platelets express a natural ligand for TREM-1. Receptor ligation activates the full repertoire of effector functions. TREM-1 is also produced in a soluble form (sTREM-1) of 17 kDa which is released in humans after endotoxin exposition or in patients suffering from severe pneumonia or sepsis. sTREM-1 can be measured in biological fluids and may be useful as a diagnostic tool.

**Synonyms:**

TREM-1