

Product datasheet for **AM26260PU-N**

Complement C5 (C5) (N-term) Mouse Monoclonal Antibody [Clone ID: 561]

Product data:

Product Type:	Primary Antibodies
Clone Name:	561
Applications:	ELISA, FN
Recommended Dilution:	Functional Assays (Ref.1,2). Immunoassays (Ref.2): Can be used as Detector in ELISA. The typical starting working dilution is 1/50.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Human C5/C5a
Specificity:	The Monoclonal antibody 561 recognizes the N-terminus of C5. The antibody reacts both with intact C5 (190 kDa) as with C5a (115 kDa).
Formulation:	PBS State: Purified State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 0.1% BSA
Concentration:	lot specific
Purification:	Protein G Chromatography
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C. DO NOT FREEZE!
Stability:	Shelf life: one year from despatch.
Gene Name:	complement component 5
Database Link:	Entrez Gene 727 Human P01031



[View online »](#)

Background:

The complement system is composed of over 30 proteins, activated in response to tissue injury, invading pathogens or other foreign surfaces. The complement pathways can be divided in the activation pathways and lytic pathway. The activation pathways lead via C3 to the cleavage of the fifth complement component C5 into C5a and C5b, resulting in activation of the lytic pathway

C5a was first described as a cleavage product of C5 with chemotactic and anaphylatoxic properties. Further characterization revealed that C5a is an essential part of the innate immune response and evidence now suggests that it may also play a role in adaptive immunity. Complement fragment C5a is a 74 residue pro-inflammatory polypeptide. C5a induces smooth muscle contraction, increases vascular permeability, causes degranulation of mast cells and basophils, and release of lysosomal enzymes. In addition C5a stimulates the directed migration of neutrophils, eosinophils, basophils and monocytes. C5a binds to at least two seven-transmembrane domain receptors, C5aR (C5R1, CD88) and C5L2 (gpr77), expressed ubiquitously on a wide variety of cells but particularly on the surface of immune cells like macrophages, neutrophils and T cells. The former is a well-established receptor that initiates G-protein-coupled signaling via mitogen-activated protein kinase pathways, thereby by inducing synthesis of cytokines such as TNF-alpha, IL-1beta, IL-6 and IL-8. Its in vivo blockade greatly reduces inflammatory injury. Much less is known about C5L2, occupancy of which by C5a does not initiate increased intracellular Ca(2+). The widespread expression of C5a receptors throughout the body allows C5a to elicit a broad range of effects. Thus, C5a has been found to be a significant pathogenic driver in a number of immuno-inflammatory diseases. Nowadays C5a is also implicated in non-immunological functions associated with developmental biology, CNS development and neurodegeneration, tissue regeneration, and haematopoiesis.

Synonyms:

CPAMD4, Complement component 5