

## Product datasheet for **AM26266PU-N**

### PI3 (C-term) Mouse Monoclonal Antibody [Clone ID: TRAB2F]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	TRAB2F
Applications:	ELISA, IHC, WB
Recommended Dilution:	Immunoassay. Western blot(non-reducing conditions): The typical starting working dilution is 1:10. Immunohistochemistry on paraffin sections: The typical starting working dilution is 1:10. Not useful for Immunohistochemistry on frozen sections.
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Specificity:	TRAB2F is an Elafin/SKALP specific antibody which shows strong reaction with fully processed Elafin/SKALP (C-terminal 57 aa) and a weaker reaction with human native full-length Elafin/SKALP. The antibody can be used for purification of human native full-length Elafin/SKALP.
Formulation:	PBS State: Purified State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 0.1% bovine serum albumin Preservative: 0.02% sodium azide
Concentration:	lot specific
Purification:	Protein G
Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C.
Stability:	Shelf life: one year from despatch.
Gene Name:	peptidase inhibitor 3
Database Link:	<a href="#">Entrez Gene 5266 Human P19957</a>



[View online »](#)

**Background:**

Elafin is an epithelial proteinase inhibitor also known under various other names such as Skin-derived Anti leukoproteinase (SKALP) and Elastase-Specific Inhibitor (ESI). Elafin belongs to the Trappin gene family and was given the systematic name Trappin-2. The Trappin family is defined by a N-terminal transglutaminase substrate domain and a C-terminal four disulphide core. Trappins have been suggested to play a role in the regulation of inflammation and in protection against tissue damage in stratified epithelia. Elafin is an inhibitor of leukocyte elastase and proteinase-3 and is a substrate for transglutaminases. The protein is constitutively expressed in various epithelia including hair follicles, oesophagus, vagina and oral cavity. Elafin is not present in normal human skin but is strongly induced during inflammation as in psoriasis and wound healing. Antibodies to Elafin can be used to evaluate the effects of treatment of psoriasis since its expression is significantly correlated with clinical scores. Antibodies to Elafin have also been successfully used to study differentiation in squamous cell carcinoma of the head-and-neck region, oesophagus and skin. It is also shown that Elafin possesses antimicrobial activity against gram-positive and gram-negative bacteria.

**Synonyms:**

Elastase-specific inhibitor, ESI, Peptidase inhibitor 3, PI-3, SKALP, WAP3, WFDC14