

Product datasheet for AM26273BT-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

liver FABP (FABP1) Mouse Monoclonal Antibody [Clone ID: L2B10]

Product data:

Product Type: Primary Antibodies

Clone Name: L2B10

Applications: ELISA, IHC, WB

Recommended Dilution: Immunohistochemistry on fozen sections: The typical starting working dilution is 1:50.

Immunoassay.

Western blot: The typical starting working dilution is 1:50.

Reactivity: Baboon, Canine, Human, Porcine, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Specificity: The monoclonal antibody L2B10 recognizes human liver fatty acid binding protein (L-FABP) of

both natural and recombinant origin.

Furthermore, the antibody can be used for the purification of human L-FABP.

Weak reactivity with Human H-FABP.

Formulation: PBS

Label: Biotin

State: Liquid 0.2 µm filtered lg fraction Stabilizer: 0.1% bovine serum albumin Preservative: 0.02% sodium azide

Concentration: lot specific

Purification: Protein G

Conjugation: Biotin

Storage: Store at 2 - 8 °C.

Stability: Shelf life: one year from despatch.

Gene Name: fatty acid binding protein 1

Database Link: Entrez Gene 2168 Human

P07148





liver FABP (FABP1) Mouse Monoclonal Antibody [Clone ID: L2B10] - AM26273BT-N

Background:

The L-FABP protein is derived from the human FABP1 gene. FABPs are small intracellular proteins (~13-14 kDa) with a high degree of tissue specificity that bind long chain fatty acids. They are abundantly present in various cell types and play an important role in the intracellular utilization of fatty acids, transport and metabolism. There are at least nine distinct types of FABP, each showing a specific pattern of tissue expression. Due to its small size, FABP leaks rapidly out of ischemically damaged necrotic cells leading to a rise in serum levels. Ischemically damaged tissues are characterized histologically by absence (or low presence) of FABP facilitating recognition of such areas. L-FABP is localized in the liver, kidney and intestinal epithelium.

Synonyms:

FABPL, L-FABP, Liver-type fatty acid-binding protein