

## Product datasheet for **AM26284PU-N**

### CD61 (ITGB3) Mouse Monoclonal Antibody [Clone ID: BV4]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	BV4
Applications:	ELISA, FN, IHC, IP, WB
Recommended Dilution:	<p><b>Immunohistochemistry on Frozen Sections:</b> Tissue sections were fixed in acetone. As negative control an irrelevant Mouse IgG was used (Ref.2). The typical starting working dilution is 1/50.</p> <p><b>Immunohistochemistry on Paraffin Sections:</b> Tissue sections were deparaffinized in xylene and pretreated with 10mmol/l sodium citrate buffer, pH 6.0 (Ref.3). The typical starting working dilution is 1/50.</p> <p><b>Functional Assays:</b> Antibody clone BV4 inhibits the downstream activation, reducing the mitogenic effects of two of its ligands, VEGFR-2 and HIV-1-Tat (Ref.1).</p> <p><b>Immunoassays.</b></p> <p><b>Immunoprecipitation:</b> Antibody clone BV4 precipitates beta3 integrin complexes from endothelial cell lysate.</p> <p><b>Western blot:</b> The typical starting working dilution is 1/50 (Ref.4).</p> <p><b>Positive Control:</b> Endometrium.</p>
Reactivity:	Bovine, Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Specificity:	The monoclonal antibody BV4 recognizes Human beta3 integrin subunit present in Platelet glycoprotein GPIIb-IIIa (integrin alphaIIb/beta3, CD41/CD61) and in the vitronectin receptor (integrin alphaV/beta3, CD51/CD61).
Formulation:	<p>PBS</p> <p>State: Purified</p> <p>State: Liquid 0.2 µm filtered Ig fraction</p> <p>Stabilizer: 0.1% BSA</p> <p>Preservative: 0.02% Sodium Azide</p>
Concentration:	lot specific
Purification:	Protein G Chromatography



[View online »](#)

Conjugation:	Unconjugated
Storage:	Store undiluted at 2-8°C. <b>DO NOT FREEZE!</b>
Stability:	Shelf life: one year from despatch.
Gene Name:	integrin subunit beta 3
Database Link:	<a href="#">Entrez Gene 3690 Human P05106</a>
Background:	<p>Integrins are a family of heterodimeric membrane glycoproteins expressed on diverse cell types which function as the major receptors for extracellular matrix and as cell-cell adhesion molecules. As adhesion molecules they play an important role in numerous biological processes such as platelet aggregation, inflammation, immune function, wound healing, tumour metastasis and tissue migration during embryogenesis. In addition integrins are involved in signaling pathways, transmitting signals both into and out from cells. All integrins consist of two non-covalently associated subunits, alpha and beta. At least 12 different alpha subunits and 8 beta subunits have been identified. The beta subunits all contain 56 conserved cysteines (except beta4 which has 48) which are arranged in four repeating units. The beta3 subunit is a 93kDa protein that contains a large loop in the N-terminus stabilized by intrachain disulphide bonding with the first cysteine-rich repeat.</p> <p>Platelet glycoprotein GPIIb-IIIa is expressed on platelets and megakaryoblasts. It is constitutively expressed and becomes activated on triggered platelets. Platelet glycoprotein GPIIb-IIIa binds to fibrinogen, fibronectin, vWF, vitronectin and thrombospondin. Next to this it is also a receptor for several soluble adhesive proteins. Vitronectin receptor is expressed on endothelial cells, some B cells, monocytes/macrophages, platelets and tumour cells. Vitronectin receptor binds next to vitronectin to fibrinogen, vWF, thrombospondin, fibronectin, osteopontin and collagen. Defects in human beta3 integrin are a cause of Glanzmann thrombasthenia, which is an autosomal recessive disorder characterized by mucocutaneous bleeding and the inability of this integrin to recognize macromolecular or synthetic peptide ligands.</p>
Synonyms:	Integrin beta-3, GP3A, GPIIIa