

## Product datasheet for **AM26291PU-N**

### **FABP3 Mouse Monoclonal Antibody [Clone ID: 67D3]**

#### **Product data:**

<b>Product Type:</b>	Primary Antibodies
<b>Clone Name:</b>	67D3
<b>Applications:</b>	ELISA, IP, WB
<b>Recommended Dilution:</b>	Immunassays (1). Western blot (2): Reduced sample treatment. The band size is ~15 kDa (Ref.2). The typical starting working dilution is 1:50. Immunoprecipitation (3). Positive control: Heart cells, recombinant human H-FABP.
<b>Reactivity:</b>	Human, Mouse, Rat
<b>Host:</b>	Mouse
<b>Isotype:</b>	IgG1
<b>Clonality:</b>	Monoclonal
<b>Immunogen:</b>	Purified Human H-FABP
<b>Specificity:</b>	The monoclonal antibody 67D3 recognizes human heart-type fatty acid-binding protein (H-FABP) of both natural and recombinant origin. Furthermore, this antibody is useful for the purification of H-FABP. Shows average cross reactivity with Human B-FABP in ELISA. Does not bind to Human A-FABP, Human I-FABP, Human L-FABP.
<b>Formulation:</b>	PBS State: Purified State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 0.1% bovine serum albumin Preservative: 0.02% sodium azide
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Protein G
<b>Conjugation:</b>	Unconjugated
<b>Storage:</b>	Store at 2 - 8 °C.
<b>Stability:</b>	Shelf life: one year from despatch.



[View online »](#)

**Gene Name:** fatty acid binding protein 3

**Database Link:** [Entrez Gene 2170 Human P05413](#)

**Background:** The H-FABP protein is derived from the human FABP3 gene. FABPs are small intracellular proteins (~13-14 kDa) with a high degree of tissue specificity that bind long chain fatty acids. They are abundantly present in various cell types and play an important role in the intracellular utilization of fatty acids, transport and metabolism. There are at least nine distinct types of FABP, each showing a specific pattern of tissue expression. Due to its small size, FABP leaks rapidly out of ischemically damaged necrotic cells leading to a rise in serum levels. Ischemically damaged tissues are characterized histologically by absence (or low presence) of FABP facilitating recognition of such areas. H-FABP is localized in the heart, skeletal and smooth muscle, mammary epithelial cells, aorta, distal tubules of the kidney, lung, brain, placenta, and ovary.

**Synonyms:** FABP11, MDGI, H-FABP, Heart-type fatty acid-binding protein