

Product datasheet for AM26317FC-N

OriGene Technologies, Inc.9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Siglec-H Rat Monoclonal Antibody [Clone ID: 440c]

Product data:

Product Type: Primary Antibodies

Clone Name: 440c Applications: IHC

Recommended Dilution: Immunohistochemistry on frozen sections: The typical starting working dilution is 1:50.

Flow cytometry: The typical starting working dilution is 1:50.

Reactivity: Mouse **Host:** Rat

Isotype: IgG2b

Clonality: Monoclonal

Specificity: The monoclonal antibody 440c reacts with Siglec-H, a cell-surface receptor molecule

selectively expressed on murine natural interferon producing cells (IPC), also called mouse plasmacytoid dendritic cells (pDC). It exclusively recognizes mouse Siglec-H in all lymphoid organs under both normal and inflammatory conditions. Siglec-H is normal present in the T cell zone of lymph nodes and spleen. Incubation of IPC with antibody 440c in vitro or

administration of antibody 440c in vivo reduces secretion of IFN-alpha in response to CpG

DNA without causing IPC depletion.

Formulation: PBS

Label: FITC

State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 1% bovine serum albumin Preservative: 0.02% sodium azide

Concentration: lot specific **Purification:** Protein G

Conjugation: FITC

Storage: Store at 2 - 8 °C.

Stability: Shelf life: one year from despatch.

Gene Name: sialic acid binding Ig-like lectin H





Siglec-H Rat Monoclonal Antibody [Clone ID: 440c] - AM26317FC-N

Database Link: Entrez Gene 233274 Mouse

Q3Y597

Background: Siglec H is unique among Sialic acid-binding Ig-like lectins (Siglecs) proteins because it

associates with the adaptor protein DAP12. DAP12 recognize certain viruses and CpG-DNA through TLR9, resulting in secretion of IFN-alpha, IL-12 and proinflammatory chemokines. Together these cytokines and chemokines recruite and activate NK cells and T cells as well as modulating the antigen presenting function of dendritic cells (DC). IPC themselves also

function as antigen presenting cells that expand memory T cells and induce Th1

differentiation. Therefore IPC may provide a first line of host defense against viral infections

by activating both innate and adaptive responses in vivo.

Synonyms: 6430529G09Rik; Siglec-H