

Product datasheet for **AM26334PU-N**

Mbl1 Rat Monoclonal Antibody [Clone ID: 2B4]

Product data:

Product Type:	Primary Antibodies
Clone Name:	2B4
Applications:	ELISA, WB
Recommended Dilution:	Immunoassays. Western blot (non-reduced): The typical starting working dilution is 1:10. It is recommended that solutions with a calcium concentration of 1 mM are used (2B4 is a calcium-dependent antibody). Not useful for immunohistology.
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG
Clonality:	Monoclonal
Specificity:	This antibody detects MBL-A.
Formulation:	PBS State: Purified State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 0.1% bovine serum albumin Preservative: 0.02% sodium azide
Concentration:	lot specific
Purification:	Protein G
Conjugation:	Unconjugated
Storage:	Store at 2 - 8 °C.
Stability:	Shelf life: one year from despatch.
Gene Name:	mannose-binding lectin (protein A) 1
Database Link:	Entrez Gene 17194 Mouse P39039


[View online »](#)

Background:

Mannose Binding Lectin (MBL) also called mannose- or mannan-binding protein (MBP) is a member of the group of collectins. MBL is an oligomeric lectin that recognizes carbohydrates as mannose and N-acetylglucosamine on pathogens. MBL contains a cysteine rich, a collagen like and a carbohydrate recognition domain. It forms a complex with C1r/C1s like serine proteases designated MASPs that proteolytically cleave C4, C2 and C3. MBL is able to activate the complement pathway independent of the classical and alternative complement activation pathways. The MBL-MASP pathway (better known as the lectin pathway) is antibody and C1q-independent. MBL exhibits complement-dependent antibacterial activity and acts directly as an opsonic and therefore plays an important role in innate immunity.

MBL is synthesized by hepatocytes and has been isolated from the liver or serum of several vertebrate species. Only one form of human MBL has been characterized, while two forms are found in rhesus monkeys, rabbits, rats and mice. The mouse forms are known as MBL-A and MBL-C.

The MBL-C concentrations in serum are about 6-fold compared to that of MBL-A. MBL-A, but not MBL-C was found to be an acute phase protein in casein and LPS-injection models. MBL-C exists in higher oligomeric forms than MBL-A.

Synonyms:

MBL-A; MBP-A; Mbpa; Mlb1; S-MBP