

# Product datasheet for AM26344BT-N

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Pecam1 Rat Monoclonal Antibody [Clone ID: MEC7.46]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: MEC7.46
Applications: IF, IHC, IP

**Recommended Dilution:** Immunohistochemistry on frozen sections (1,3): Tissue sections were fixed in acetone for 10

min at RT and pretreated with 0.01% hydrogen peroxide to quench endogenous peroxidases.

Normal rabbit serum was used as blocking agent. As (Ref.1). The typical starting working dilution is 1:50.

Flow cytometry (1): Stains the extracellular domain of mouse PECAM-1; 16 cells were used per sample. No fixation or permeabilization needed. Epitope is sensitive to trypsin! (Ref.1).

The typical starting working dilution is 1:50.

Immunoflourescence (1,2).

Immunoprecipitation (1): Lysate of 35S-Methionine labeled 5HV cells was used. Molecule of

~130 kDa.

Positive control: Mouse t-end.1 cells. Negative control: L929 cells (Fibroblasts).

Reactivity: Mouse
Host: Rat
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Mouse t.end.1 cells (polyoma middle T (PmT)-transformed EC)

**Specificity:** The monoclonal antibody MEC7.46 recognizes the mouse form of the platelet-endothelial cell

adhesion molecule (PECAM)-1 (CD31).

The reactivity of the monoclonal antibody MEC7.46 is restricted to the isoform of the molecule that is selectively expressed by endothelial cells. The antibody precipitates a 130 kDa molecule present on the membrane of endothelial cells of all mouse blood vessels both in normal, inflamed and tumor tissues. The antigen is predominantly present at the lateral borders of endothelial cells as described for human PECAM-1. Staining of MEC7.46 can be

seen on capillaries, veins, arteries and liver sinusoids.





#### Pecam1 Rat Monoclonal Antibody [Clone ID: MEC7.46] - AM26344BT-N

Formulation: PBS

Label: Biotin

State: Liquid 0.2 µm filtered Ig fraction Stabilizer: 0.1% bovine serum albumin Preservative: 0.02% sodium azide

Concentration:lot specificPurification:Protein GConjugation:Biotin

Storage: Store at 2 - 8 °C.

**Stability:** Shelf life: one year from despatch.

**Gene Name:** platelet/endothelial cell adhesion molecule 1

Database Link: Entrez Gene 18613 Mouse

Q08481

**Background:** PECAM-1 is a member of the immunoglobulin superfamily. This heavily glycosylated protein is

found in the entire vascular endothelium of adult mice and functions in mediating cellular adhesion by heterophilic and homophilic mechanisms. PECAM-1 is detected within the lymphopoietic islands in the spleen of newborn (day 12) and in the bone marrow of adult

mice. Capillary endothelial cells of adult mice also express PECAM-1.

Synonyms: PECAM-1, EndoCAM, GPIIA'