

Product datasheet for **AM26506AF-N**

Cd300a Rat Monoclonal Antibody [Clone ID: TX40]

Product data:

Product Type:	Primary Antibodies
Clone Name:	TX40
Applications:	FC
Recommended Dilution:	Flow cytometry: 5-10 µg/ml (final concentration). For details see protocol below.
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Mouse MAIR-I transfected Ba/F3 cells
Specificity:	This antibody reacts with mouse CD300a.
Formulation:	PBS containing 50% glycerol, pH 7.2. No preservative is contained. State: Azide Free State: Liquid Ig fraction
Concentration:	lot specific
Purification:	Protein G agarose
Conjugation:	Unconjugated
Storage:	Upon receipt, store (in aliquots) at -20 °C. Avoid repeated freezing and thawing.
Stability:	Shelf life: One year from despatch.
Gene Name:	CD300A molecule
Database Link:	Entrez Gene 217303 Mouse Q6SIQ0



[View online »](#)

Background:

Immune responses are regulated by opposing positive and negative signals triggered by the interaction of activating and inhibitory cell surface receptors with their ligands. Shibuya et al. identified novel paired activated and inhibitory immunoglobulin-like receptors, designated myeloid-associated immunoglobulin-like receptor (MAIR) I and MAIR-II, whose extracellular domains are highly conserved by each other. MAIR-I, expressed on the majority of myeloid cells, including macrophages, granulocytes, mast cells, and dendritic cells, contains the tyrosine-based sorting motif and the immunoreceptor tyrosine-based inhibitory motif-like sequences in the cytoplasmic domains. On the other hand, MAIR-II, expressed on subsets of peritoneal macrophages and B cells, associates with the immunoreceptor tyrosine-based activation motif-bearing adaptor DAP12. MAIR-I is also known as CD300a/ CMRF-35-like Ig-like molecule-8 (CLM-8)/leukocyte mono-Ig-like receptor 1 (LMIR1). MAIR-II is also known as CD300d/LMIR2/CLM-4/dendritic cell-derived Ig-like receptor 1 (Dlgr1).

Synonyms:

CLM-8, CMRF35-H, CMRF35-H9, CMRF-35-H9, IRC1/IRC2, IRp60

Note:

This product was originally produced by MBL International.

Protocol:
Flow cytometric analysis for floating cells

We usually use Fisher tubes or equivalents as reaction tubes for all steps described below.

- 1) Wash the cells 3 times with washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.1% NaN₃].
- 2) Resuspend the cells with washing buffer (5 x 10⁶ cells/mL).
- 3) Add 50 µL of the cell suspension into each tube, and centrifuge at 500 x g for 1 minute at room temperature (20~25°C). Remove supernatant by careful aspiration.
- 4) Add 10 µL of normal goat serum containing 1 mg/mL normal human IgG and 0.1% NaN₃ to the cell pellet after tapping. Mix well and incubate for 5 minutes at room temperature.
- 5) Add 40 µL of the primary antibody at the concentration as suggest in the APPLICATIONS diluted in the washing buffer. Mix well and incubate for 30 minutes at room temperature.
- 6) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 7) Add 30 µL of 1:100 PE conjugated anti-rat IgG diluted with the washing buffer. Mix well and incubate for 15 minutes at room temperature.
- 8) Add 1 mL of the washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 9) Resuspend the cells with 500 µL of the washing buffer and analyze by a flow cytometer. (Positive control for Flow cytometry; WEHI-3B)


Flow cytometric analysis for whole blood cells

We usually use Falcon tubes or equivalents as reaction tubes for all steps described below.

- 1) Add 50 µL of mouse CD300a monoclonal antibody (TX40) at the concentration as suggest in the APPLICATIONS diluted in the washing buffer [PBS containing 2% fetal calf serum (FCS) and 0.1% NaN₃] into each tube.
- 2) Add 50 µL of whole blood into each tube. Mix well, and incubate for 30 minutes at room temperature (20~25 °C).

- 3) Add 1 mL of washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 4) Add 30 μ L of 1:100 PE conjugated anti-rat IgG diluted with washing buffer. Mix well and incubate for 15 minutes at room temperature.
- 5) Add 1 mL of washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 6) Lyse with OptiLyse C (for analysis on Beckman Coulter instruments) or OptiLyse B (for analysis on BD instruments), using the procedure recommended in the respective package inserts.
- 7) Add 1 mL of H₂O to each tube and incubate for 10 minutes at room temperature.
- 8) Centrifuge at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 9) Add 1 mL of washing buffer followed by centrifugation at 500 x g for 1 minute at room temperature. Remove supernatant by careful aspiration.
- 10) Resuspend the cells with 500 μ L of the washing buffer and analyze by a flow cytometer.

Product images:



Flow cytometric analysis of mouse CD300a expression on WEHI-3B cells. Open histogram indicates the reaction of isotypic control to the cells. Shaded histogram indicates the reaction of AM26506AF-N to the cells.