

Product datasheet for AM26741PU-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CBX3 Mouse Monoclonal Antibody [Clone ID: 2MOD-1G6]

Product data:

Product Type: Primary Antibodies

Clone Name: 2MOD-1G6

Applications: IHC

Recommended Dilution: Immunohistochemistry on paraffin sections: 0.4 µg/ml (1/500).

Suggested positive control: Human / Porcine brain cortex.

Reactivity: Human, Mouse, Porcine

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Recombinant mouse HP1y

Specificity: This antibody detects HP1y.

Formulation: Phosphate buffered saline pH 7.2 (PBS)

State: Purified

State: Lyophilized purified Ig fraction

Stabilizer: 5 mg/ml bovine serum albumin (BSA)

Preservative: 0.01 % Kathon

Reconstitution Method: Reconstitute by adding 0.5 ml distilled water. This stock solution contains 0.2 mg/ml lgG.

Conjugation: Unconjugated

Storage: Store lyophilized at 2-8°C for 6 month or at -20°C long term.

After reconstitution store the antibody undiluted at 2-8°C for one month

or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Database Link: P23198






Background:

Heterochromatin Protein 1 (HP1) ("Chromobox Homolog", CBX) includes three isoforms in mammals, designated α , β , and γ . HP1 proteins are approximately 25 kDa in size. They are highly conserved adapter proteins with important regulatory functions in the cell nucleus. These functions include notably gene silencing by heterochromatin formation. HP1 proteins are fundamental units of heterochromatin packaging that are enriched at the centromeres and telomeres of nearly all Eukaryotic chromosomes. HP1 β and γ also localize to euchromatic sites in the genome. Members of the HP1 family are characterized by an N-terminal chromodomain and a C-terminal chromoshadow domain, separated by a Hinge region. Chromodomain and chromoshadow domain establish secondary interactions with a large number of other proteins. HP1 proteins are subject to multiple types of post-translational modifications, including phosphorylation, acetylation, methylation, ubiquitination and sumoylation, suggesting multiple means of regulation.


Synonyms:

Chromobox protein homolog 3, HECH, HP1 gamma, HP1-gamma, HP1Hs-gamma

Product images:



Immunohistochemistry on paraffin sections: Human brain cortex, stained withAM26741PU-N.



Immunohistochemistry on paraffin sections: Swine brain cortex, stained withAM26741PU-N.