

Product datasheet for AM26784AC-N

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD94 (KLRD1) Mouse Monoclonal Antibody [Clone ID: HP-3D9]

Product data:

Product Type: Primary Antibodies

Clone Name: HP-3D9

Applications: FC

Recommended Dilution: Flow Cytometry analysis of human blood cells using 10 μl reagent / 100 μl of whole blood

or 106 cells in a suspension.

The content of a vial (1 ml) is sufficient for 100 tests.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Cultured human NK cells

Specificity: This antibody recognizes CD94, a 70 kDa type II transmembrane glycoprotein expressed on

NK cells, NK-T cells, and subsets of CD8+ T cells and gamma/delta T cells.

Formulation: Phosphate buffered saline (PBS)

Label: APC

State: Liquid purified Ig fraction

Stabilizer: 0.2% (w/v) high-grade protease free BSA

Preservative: 15 mM sodium azide

Label: Conjugated with cross-linked Allophycocyanin (APC) under optimum conditions. The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No

reconstitution is necessary.

Purification: The conjugate is purified by size-exclusion chromatography and adjusted for direct use. No

reconstitution is necessary.

Conjugation: APC

Storage: Store the antibody undiluted at 2-8°C.

DO NOT FREEZE!

This product is photosensitive and should be protected from light.

Stability: Shelf life: one year from despatch.

Gene Name: killer cell lectin like receptor D1





Database Link: Entrez Gene 3824 Human

Q13241

Background: CD94, also known as KLRD1 (killer cell lectin-like receptor D1), is a transmembrane

glycoprotein of the C-type lectin family, which forms disulfide-linked heterodimers with NKG2A, B, C, E, H proteins, constituting functionally distinct receptors of NK cells and related


cell types. CD94/NKG2A and CD94/NKG2B heterodimers serve as inhibitory, whereas

CD94/NKG2C and CD94/NKG2E as activating receptors. The ligand for CD94/NKG2 complexes has been identified as HLA-E. Extent of CD94 expression on NK cell surface can be used to

demonstrate their progress through the differentiation process.

Synonyms: KLRD1, KP43, NK cell receptor

Product images:



Surface staining of CD94 in human peripheral blood with anti-CD94 (HP-3D9) APC.