

## **Product datasheet for AM26784PU-N**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### CD94 (KLRD1) Mouse Monoclonal Antibody [Clone ID: HP-3D9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: HP-3D9

Applications: FC

Recommended Dilution: Flow cytometry.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Cultured human NK cells

**Specificity:** This antibody recognizes CD94, a 70 kDa type II transmembrane glycoprotein expressed on

NK cells, NK-T cells, and subsets of CD8+T cells and gamma/delta T cells.

Formulation: Phosphate buffered saline (PBS)

State: Purified

State: Liquid Ig fraction

Preservative: 15 mM sodium azide, approx. pH 7.4

**Concentration:** lot specific

**Purification:** Protein-A affinity chromatography (> 95% pure by SDS-PAGE)

Conjugation: Unconjugated

Storage: Store undiluted at 2-8°C.

**DO NOT FREEZE!** 

Stability: Shelf life: one year from despatch.

Gene Name: killer cell lectin like receptor D1

Database Link: Entrez Gene 3824 Human

Q13241




Background:

CD94, also known as KLRD1 (killer cell lectin-like receptor D1), is a transmembrane glycoprotein of the C-type lectin family, which forms disulfide-linked heterodimers with NKG2A, B, C, E, H proteins, constituting functionally distinct receptors of NK cells and related cell types. CD94/NKG2A and CD94/NKG2B heterodimers serve as inhibitory, whereas CD94/NKG2C and CD94/NKG2E as activating receptors. The ligand for CD94/NKG2 complexes has been identified as HLA-E. Extent of CD94 expression on NK cell surface can be used to demonstrate their progress through the differentiation process.

Synonyms: KLRD1, KP43, NK cell receptor

# **Product images:**



Surface staining of CD94 in human peripheral blood with anti-CD94 (HP-3D9) purified, GAM-