

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM31025PU-N

SAPK4 (MAPK13) Mouse Monoclonal Antibody [Clone ID: 2B2]

Product data:


Product Type:	Primary Antibodies
Clone Name:	2B2
Applications:	ELISA, IF, IHC, WB
Recommended Dilution:	ELISA. Immunofluorescence: 10 μg/ml. Immunohistochemistry on Paraffin Sections: 5 μg/ml. Western Blot.
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	MAPK13 antibody was raised against recombinant protein
Specificity:	This antibody reacts to MAPK13 / SAPK4.
Formulation:	PBS, pH 7.2 State: Purified State: Liquid purfied lg fraction
Concentration:	lot specific
Purification:	Protein A chromatography
Conjugation:	Unconjugated
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	mitogen-activated protein kinase 13
Database Link:	<u>Entrez Gene 5603 Human</u> <u>O15264</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	SAPK4 (MAPK13) Mouse Monoclonal Antibody [Clone ID: 2B2] – AM31025PU-N
Background:	MAP kinase p38 delta, a MAPK type protein kinase, is an environmental stress- and inflammatory cytokine-induced kinase that is associated with the regulation of gene expression and cell proliferation. p38delta is phosphorylated by MEK3/MAP2K3 or MKK6/MAP2K6. Activated p38delta has been shown to phosphorylate ATF2, eEF2K, MBP and PHAS-1.
Synonyms:	Mitogen-activated protein kinase 13, Stress-activated protein kinase 4, p38 delta, PRKM13, SAPK4
Protein Families:	Druggable Genome, Protein Kinase
Protein Pathway	S: Amyotrophic lateral sclerosis (ALS), Epithelial cell signaling in Helicobacter pylori infection, Fc epsilon RI signaling pathway, GnRH signaling pathway, Leukocyte transendothelial migration, MAPK signaling pathway, Neurotrophin signaling pathway, NOD-like receptor signaling pathway, Progesterone-mediated oocyte maturation, RIG-I-like receptor signaling pathway, T cell receptor signaling pathway, Toll-like receptor signaling pathway, VEGF signaling pathway

Product images:




MAPK13 monoclonal antibody clone 2B2 Western Blot analysis of MAPK13 expression in A-431.




Detection limit for recombinant GST tagged MAPK13 is approximately 0.03ng/ml as a capture antibody.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Human Prostate: Formalin-Fixed, Paraffin-Embedded (FFPE)



Immunofluorescence of monoclonal antibody to MAPK13 on A-431 cell. [antibody concentration 10 ug/ml]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US