

Product datasheet for AM31739PU-N

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

CREB1 Mouse Monoclonal Antibody [Clone ID: 3F1-1B2]

Product data:

Product Type: Primary Antibodies

Clone Name: 3F1-1B2

Applications: ELISA, IF, IHC, WB

Recommended Dilution: ELISA.

Immunofluorescence: 10 µg/ml.

Immunohistochemistry on Paraffin Sections: 5 µg/ml.

Western Blot: 1/500 - 1/1000.

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: CREB1 antibody was raised against cREB1 (AAH10636, 1 a.a. ~ 342 a.a) full length

recombinant protein with GST tag.

Specificity: Recognizes CREB1

Formulation: PBS, pH 7.2

State: Purified

State: Liquid purified Ig fraction

Purification: Protein A Chromatography

Conjugation: Unconjugated

Storage: Store the antibody (in aliquots) at -20°C or -80°C.

Avoid freeze-thaw cycles.

Stability: Shelf life: one year from despatch.

Gene Name: cAMP responsive element binding protein 1

Database Link: Entrez Gene 1385 Human

P16220



Background:


CREB (cAMP response element binding protein1), a DNA binding protein, is a member of the leucine zipper family of transcription factors. CREB interaction with DNA requires phosphorylation at Ser 119 by activated cAMP protein kinase A. Once phosphorylated CREB binds to an 8 base enhancer know as the cAMP response element (CRE) containing a conserved core sequence TGACG. CREB also interacts with CBP (CREB binding protein) and RNA polymerase II when phosphorylated on Ser 133. Different isoforms of this protein are the result of two transcript variants.

Synonyms: CREB-1


Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Antigen processing and presentation, Huntington's disease, Melanogenesis, Prostate cancer

Product images:




Western Blot analysis of CREB1 expression in transfected 293T cell line. Lane 1: CREB1 transfected lysate (37 KDa). Lane 2: Nontransfected lysate.




Detection limit for recombinant GST tagged CREB1 is approximately 1ng/ml as a capture antibody.






Human Breast: Formalin-Fixed, Paraffin-Embedded (FFPE)



Human Tonsil: Formalin-Fixed, Paraffin-Embedded (FFPE)



Immunofluorescence of monoclonal antibody to CREB1 on HeLa cell at 10 ug/ml.