

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for AM31856RP-L

BCL3 Hamster Monoclonal Antibody [Clone ID: Ham150-3.5]

Product data:

| Product Type: | Primary Antibodies |
|-----------------------|---|
| Clone Name: | Ham150-3.5 |
| Applications: | FC |
| Recommended Dilution: | This Clone has been described to work in ELISA , Western Blot and Flow Cytometry. |
| Reactivity: | Human, Mouse |
| Host: | Hamster |
| lsotype: | lgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Recombinant Bcl-3-6-histidine tag from Armenian Hamster. Fusion Partner: SP2/0 myeloma. |
| Specificity: | Recognizes B-cell Leukemia/lymphoma 3 |
| Formulation: | PBS containing 0.02% Sodium Azide as preservative and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: PE State: Liquid purified IgG fraction. |
| Concentration: | lot specific |
| Purification: | Protein G Affinity Chromatography. |
| Conjugation: | PE |
| Storage: | Store the antibody undiluted at 2-8°C. DO NOT FREEZE! Avoid prolonged exposure to light. |
| Stability: | Shelf life: one year from despatch. |
| Gene Name: | B-cell CLL/lymphoma 3 |
| Database Link: | <u>Entrez Gene 602 Human</u> <u>P20749</u> |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | BCL3 Hamster Monoclonal Antibody [Clone ID: Ham150-3.5] – AM31856RP-L |
|-------------------------------|--|
| Background: | B cell Lymphoma 3 is a member of the IkB subfamily of inhibitors. This subfamily is part of the NF-kB transcription factor protein family and it is suggested that a balance in the concentration of various NF-kB family members regulates apoptosis and survival of activated T cells. |
| | Bcl-3 has been shown to have versatile functions such as cytoplasmic activation of p50 homodimers, translocation to the nucleus, and modulation of the transcriptional machinery in the nucleus. Bcl-3 activity often relies on several nuclear interacting proteins such as Tip60, Jab1, Bard1, and Pirin. The Mouse Bcl-3 coding region exhibits 80% homology with Human Bcl-3 which is associated with human B-cell chronic lymphocytic leukemias (CLLs). Bcl-3 is detected in various tissues such as spleen and other lymphoid organs. |
| Synonyms: Protein Families | BCL3, BCL4, D19S37 Druggable Genome, Transcription Factors |
| | |

~ \$%~

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US