

Product datasheet for **AM31862AC-N**

T Cell Receptor (TCR) V alpha-2 Rat Monoclonal Antibody [Clone ID: B20.1]

Product data:

Product Type:	Primary Antibodies
Clone Name:	B20.1
Applications:	FC
Recommended Dilution:	Flow Cytometry (See Protocols). This clone has also been reported to work in Immunoprecipitation . (1,2)
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2a
Clonality:	Monoclonal
Specificity:	This antibody reacts with Mouse T-Cell Receptor (TCR) Va2 chains (1), and recognizes the majority of the TCR Va2 subfamily in mice carrying the a, b and c haplotypes 1,2. It also reacts with the products of T cell receptor, Vδ8 due to the high degree of homology (1).
Formulation:	PBS containing 0.09% Sodium Azide as preservative and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: APC State: Liquid purified IgG fraction.
Concentration:	lot specific
Purification:	Protein G Affinity Chromatography.
Conjugation:	APC
Storage:	Store the antibody undiluted at 2-8°C. DO NOT FREEZE! Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Background:	The TCR alpha chain complexes with the TCR beta chain to form the T cell receptor in 95% of T cells, whereas the remaining 5% of T cells express gamma and delta chains (γ/δ). TCR Va2 is a distinct TCR subfamily found in mice having the a, b, and c haplotypes.



[View online »](#)

Note:

Protocol: **Flow Cytometry Analysis:**

NOTE: Preblocking of Fc receptors for 10 minutes using 0.5 µg of purified anti-Mouse CD16/32 is recommended.

Method:

1. Prepare cell suspension in Media A. For cell replications, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration 2×10^7 cells/ml in media A. Add 50 µl of this suspension to each tube (each tube will then contain 1×10^6 cells, representing one test).
4. To each tube add 0.25 µg of this antibody AM31862AC-N per 1×10^6 cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate tubes at 4°C for 30-60 minutes. (It is recommended that the tubes are protected from light since most fluorochromes are light sensitive).
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50 µl ice cold Media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15 µl of propidium iodide at 0.5 mg/ml in phosphate buffered saline. This stains dead cells by intercalating DNA.

Media:

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100 µl of 2 M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% bovine serum albumin + sodium azide (100 µl of 2 M sodium azide in 100 mls).