

## Product datasheet for **AM31873RP-N**

### CD45 / LCA (CD45RC) Rat Monoclonal Antibody [Clone ID: IBL-8]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	IBL-8
Applications:	FC
Recommended Dilution:	Immunohistochemistry on acetone-fixed frozen sections. Flow Cytometry. Immunoprecipitation.
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Mouse Spleen Cells <u>Donor:</u> Wistar spleen <u>Fusion Partner:</u> Sp-2/0 Ag14
Specificity:	Anti-mouse CD45RC is against the exon C-dependent RC isoform and reacts strongly with B cells, and less intensely with most CD8+ T cells. It does not recognize CD4+ T cells. Also, myeloid cells do not express the RC isoform.
Formulation:	PBS containing 0.02% sodium azide (NaN <sub>3</sub> ) as preservative and EIA grade BSA as a stabilizing protein to bring total protein concentration to 4-5 mg/ml. Label: PE State: Liquid purified Ig fraction Label: R - Phycoerythrin
Concentration:	lot specific
Purification:	Affinity purified
Conjugation:	PE
Storage:	Store the antibody undiluted at 2-8°C. <b>DO NOT FREEZE!</b>
Stability:	Shelf life: one year from despatch.
Gene Name:	protein tyrosine phosphatase, receptor type, C



[View online »](#)

**Database Link:** [Entrez Gene 19264 Mouse P06800](#)

**Background:** CD45 (L-CA) is a transmembrane phosphotyrosine phosphatase expressed on leukocytes.

**Synonyms:** PTPRC, Leukocyte common antigen, L-CA, T200

**Note:** Protocol: **FLOW CYTOMETRY ANALYSIS:**

**Method:**

1. Prepare a cell suspension in media A. For cell preparations, deplete the red blood cell population with Lympholyte®-M cell separation medium.
2. Wash 2 times.
3. Resuspend the cells to a concentration of  $2 \times 10^7$  cells/ml in media A. Add 50  $\mu$ l of this suspension to each tube (each tube will then contain  $1 \times 10^6$  cells, representing 1 test).
4. To each tube, add 1.0  $\mu$ g of this antibody per  $10^6$  cells.
5. Vortex the tubes to ensure thorough mixing of antibody and cells.
6. Incubate the tubes for 30 minutes at 4°C.  
(It is recommended that the tubes are protected from light, since most fluorochromes are light sensitive.)
7. Wash 2 times at 4°C.
8. Resuspend the cell pellet in 50  $\mu$ l ice cold media B.
9. Transfer to suitable tubes for flow cytometric analysis containing 15  $\mu$ l of propidium iodide at 0.5 mg/ml in PBS. This stains dead cells by intercalating in DNA.

**Media:**

- A. Phosphate buffered saline (pH 7.2) + 5% normal serum of host species + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).
- B. Phosphate buffered saline (pH 7.2) + 0.5% Bovine serum albumin + sodium azide (100  $\mu$ l of 2M sodium azide in 100 mls).

**Results:**

Tissue Distribution by Flow Cytometry Analysis:


Mouse Strain: BALB/c

Cell Concentration:  $1 \times 10^6$  cells per test

Antibody Concentration Used: 1.0  $\mu$ g/ $10^6$  cells

Isotypic Control: PE Rat IgG1

## Product images:



Percentage of cells stained above control: 34.8%