

Product datasheet for **AM31880PU-N**

Adipoq Rat Monoclonal Antibody [Clone ID: 7J61]

Product data:

Product Type:	Primary Antibodies
Clone Name:	7J61
Applications:	WB
Recommended Dilution:	Western Blot: 1/500-1/1000
Reactivity:	Mouse
Host:	Rat
Isotype:	IgG2
Clonality:	Monoclonal
Immunogen:	Purified Mouse Recombinant Adiponectin.
Specificity:	This antibody detects Mouse Adiponectin. It is crossreactive to Human Adiponectin protein in Western blotting.
Formulation:	0.2 µm filtered solution in PBS State: Purified State: Lyophilized purified IgG fraction of Culture Supernatant
Reconstitution Method:	Restore with 200 µl sterile PBS and the final concentration is 500 µg/ml
Purification:	Protein A/G Affinity Chromatography
Conjugation:	Unconjugated
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	adiponectin, C1Q and collagen domain containing
Database Link:	<u>Entrez Gene 11450 Mouse Q60994</u>



[View online »](#)

Background:

Adiponectin, also known as Acrp30, is an adipocyte-derived protein with wide ranging paracrine and endocrine effects on metabolism and inflammation. It promotes adipocyte differentiation, fatty acid catabolism, and insulin sensitivity, and is negatively correlated with obesity, type 2 diabetes, and atherogenesis. In this context, adiponectin is an antiinflammatory agent, but it exerts proinflammatory effects in nonmetabolic disorders such as rheumatoid arthritis and inflammatory bowel disease. Adiponectin interacts with the receptors AdipoR1 and AdipoR2, calreticulin, and Cadherin13/ T-Cadherin, as well as with several growth factors. Mature mouse adiponectin consists of a 66 amino acid (aa) N-terminal collagenous region and a 137 aa C terminal C1q like globular domain which can be cleaved by a leukocytederived elastase. Mature mouse adiponectin shares 83% and 91% amino acid (aa) sequence identity with human and rat adiponectin, respectively. Adiponectin associates into trimers that may assemble into medium molecular weight (MMW) hexamers and then into >300 kDa high molecular weight (HMW) oligomers. The glycosylation of four hydroxylated lysine residues in the collagenous domain is required for the intracellular formation of HMW complexes. The various multimeric forms of adiponectin exhibit distinct tissue specific and gender specific profiles and activities.

Synonyms:

ADIPOQ, ACDC, ACRP30, APM1, GBP28